


AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application.

Listing of Claims:

1. (currently amended) A compound of the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of:

(i) [[B is]] phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³², (b) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³⁶, (c) a ring carbon or nitrogen, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³², is optionally substituted by R³³, (d) a ring carbon or nitrogen, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³⁶, is optionally substituted by R³⁵, and (e) a ring carbon or nitrogen, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R³³ and R³⁵, respectively, is optionally substituted by R³⁴.

~~a nitrogen with a removable hydrogen or a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³², a nitrogen with a removable hydrogen or a carbon at the other position adjacent to the~~

~~point of attachment is optionally substituted by R³⁶, a nitrogen with a removable hydrogen or a carbon adjacent to R³² and two atoms from the point of attachment is optionally substituted by R³³, a nitrogen with a removable hydrogen or a carbon adjacent to R³⁶ and two atoms from the point of attachment is optionally substituted by R³⁵, and a nitrogen with a removable hydrogen or a carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;~~

(ii) hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B may be optionally substituted at any carbon up to 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶; and

(iii) a C3-C12 cycloalkyl or a C4-C9 heterocyclyl, wherein (a) each ring carbon may be optionally substituted with R₃₃, (b) a ring carbon, other than the ring carbon at the point of attachment of B to A, may be optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, (c) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (d) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (e) a ring carbon or nitrogen atom, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (f) a ring carbon or nitrogen, if present, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², (g) a ring carbon or nitrogen, if present, in a first gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹⁰, is optionally substituted by R¹¹, (h) a ring carbon or nitrogen, if present, in a second gamma position relative to the carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹², is optionally substituted by R³³, and (i) a ring carbon or nitrogen, if present, in a delta position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹¹ and R³³, respectively, is optionally substituted by R³⁴;

R⁹, R¹⁰, R¹¹, R¹², and R¹³, ~~R³², R³³, R³⁴, R³⁵, and R³⁶~~ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino,

alkylenedioxy, haloalkylthio, alkanoyloxy, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, alkoxyalkyl, haloalkoxylalkyl, hydroxy, amino, alkoxyamino, nitro, alkylamino, N-alkyl-N-arylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylthio, alkylthioalkyl, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, alkylsulfonylalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkylsulfonamido, amidosulfonyl, alkanoyl, haloalkanoyl, alkyl, alkenyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyhaloalkyl, hydroxyalkyl, aminoalkyl, haloalkoxyalkyl, carboxyalkyl, carboalkoxy, carboxy, carboxamido, carboxamidoalkyl, and cyano;

R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} are independently selected from the group consisting of:

(i) hydrido, acetamido, haloacetamido, amidino, guanidino, alkylenedioxy, haloalkylthio, alkanoyloxy, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, alkoxyalkyl, haloalkoxylalkyl, hydroxy, amino, alkoxyamino, nitro, alkylamino, N-alkyl-N-arylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylthio, alkylthioalkyl, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, alkylsulfonylalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkylsulfonamido, amidosulfonyl, alkanoyl, haloalkanoyl, alkyl, alkenyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyhaloalkyl, hydroxyalkyl, aminoalkyl, haloalkoxyalkyl, carboxyalkyl, carboalkoxy, carboxy, carboxamido, carboxamidoalkyl, and cyano; and

(ii) optionally Q^b ;

~~B is optionally selected from the group consisting of hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B may be optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} ;~~

~~B is optionally a C3-C12 cycloalkyl or a C4-C9 heterocyclyl, wherein each ring carbon may be optionally substituted with R^{33} , a ring carbon other than the ring carbon at the point of attachment of B to A may be optionally substituted with oxo provided that~~

~~no more than one ring carbon is substituted by oxo at the same time, ring carbons and nitrogens adjacent to the carbon at the point of attachment may be optionally substituted with R^9 or R^{13} , a ring carbon or nitrogen adjacent to the R^9 position and two atoms from the point of attachment may be substituted with R^{10} , a ring carbon or nitrogen adjacent to the R^{13} position and two atoms from the point of attachment may be substituted with R^{12} , a ring carbon three atoms from the point of attachment and adjacent to the R^{10} position may be substituted with R^{11} , a ring carbon three atoms from the point of attachment and adjacent to the R^{12} position may be substituted with R^{33} , and a ring carbon four atoms from the point of attachment and adjacent to the R^{11} and R^{33} positions may be substituted with R^{34} ;~~

A is selected from the group consisting of a bond, $(W^7)_{rr}-(CH(R^{15}))_{pa}$, and $(CH(R^{15}))_{pa}-(W^7)_{rr}$ wherein rr is 0 or 1, pa is an integer selected from 0 through 6, and W^7 is selected from the group consisting of O, S, C(O), $(R^7)NC(O)$, $(R^7)NC(S)$, and $N(R^7)$, with the proviso that no more than one of the group consisting of rr and pa is 0 ~~at the same time~~;

R^7 is selected from the group consisting of hydrido, hydroxy, and alkyl;

R^{15} is selected from the group consisting of hydrido, hydroxy, halo, alkyl, and haloalkyl;

Ψ is NH or NOH;

M is N or R^1-C ;

R^1 is selected from the group consisting of hydrido, alkyl, alkenyl, cyano, halo, haloalkyl, haloalkoxy, haloalkylthio, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

R^2 is Z^0-Q ;

Z^0 is selected from the group consisting of:

(i) a bond, $W^0-(CH(R^{42}))_p$ wherein p is an integer selected from 0 through 3 and W^0 is selected from the group consisting of O, S, C(O), S(O), $N(R^{41})$, and $ON(R^{41})$, $(CH(R^{41}))_g-O$ wherein g is an integer selected from 1 through 3, and $(CH(R^{41}))_g-S$ wherein g is an integer selected from 1 through 3, with the proviso that Z^0 is directly bonded to the pyrimidinone ring; **and**

(ii) ~~Z^0 is optionally~~ $W^{22}-(CH(R^{42}))_h$ wherein h is 0 or 1 and W^{22} is selected from the group consisting of $CR^{41}=CR^{42}$, 1,2-cyclopropyl, 1,2-cyclobutyl, 1,2-cyclohexyl, 1,3-cyclohexyl, 1,2-cyclopentyl, 1,3-cyclopentyl, 2,3-morpholinyl, 2,4-morpholinyl, 2,6-morpholinyl, 3,4-morpholinyl, 3,5-morpholinyl, 1,2-piperazinyl, 1,3-piperazinyl, 2,3-piperazinyl,

2,6-piperazinyl, 1,2-piperidinyl, 1,3-piperidinyl, 2,3-piperidinyl, 2,4-piperidinyl, 2,6-piperidinyl, 3,4-piperidinyl, 1,2-pyrrolidinyl, 1,3-pyrrolidinyl, 2,3-pyrrolidinyl, 2,4-pyrrolidinyl, 2,5-pyrrolidinyl, 3,4-pyrrolidinyl, 2,3-tetrahydrofuranyl, 2,4-tetrahydrofuranyl, 2,5-tetrahydrofuranyl, and 3,4-tetrahydrofuranyl, wherein Z^0 is directly bonded to the pyrimidinone ring and W^{22} is optionally substituted with one or more substituents selected from the group consisting of R^9 , R^{10} , R^{11} , R^{12} , and R^{13} ;

R^{41} and R^{42} are independently selected from the group consisting of amidino, hydroxyamino, hydrido, hydroxy, amino, and alkyl;

Q is selected from the group consisting of:

(i) [[Q is]] phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon or nitrogen, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon or nitrogen, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon or nitrogen, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11} with the proviso that Q is other than phenyl when Z^0 is a bond; ~~a nitrogen with a removable hydrogen or a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , a nitrogen with a removable hydrogen or a carbon at the other position adjacent to the point of attachment is optionally substituted by R^{13} , a nitrogen with a removable hydrogen or a carbon adjacent to R^9 and two atoms from the point of attachment is optionally substituted by R^{10} , a nitrogen with a removable hydrogen or a carbon adjacent to R^{13} and two atoms from the point of attachment is optionally substituted by R^{12} , and a nitrogen with a removable hydrogen or a carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than phenyl when Z^0 is a bond; and~~

(ii) ~~Q is optionally hydrido with the proviso that Z^0 is other than a bond;~~


K is $(CR^{4a}R^{4b})_n$ wherein n is 1 or 2;

R^{4a} and R^{4b} are independently selected from the group consisting of halo, hydrido, hydroxyalkyl, alkyl, alkoxyalkyl, alkylthioalkyl, and haloalkyl;

E^0 is E^1 , when K is $(CR^{4a}R^{4b})_n$, wherein E^1 is selected from the group consisting of a bond, C(O), C(S), C(O)N(R^7), (R^7)NC(O), S(O)₂, (R^7)NS(O)₂, and S(O)₂N(R^7);

Y^0 is selected from the group consisting of:

(i) [[Y^0 is]] the formula



wherein J^5 , J^6 , D^5 , D^6 and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J^5 and J^6 is absent when T is a 5-membered heteroaryl ring, J^5 is optionally substituted by R^{17} when J^5 is a carbon atom, J^6 is optionally substituted by R^{18} when J^6 is a carbon atom, D^5 is optionally substituted by R^{16} when D^5 is a carbon atom and D^6 is optionally substituted by R^{19} when D^6 is a carbon atom; phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s , a carbon two or three contiguous atoms from the point of attachment of Q^s to the phenyl or heteroaryl ring is substituted by Q^b , a carbon adjacent to the point of attachment of Q^s is optionally substituted by R^{17} , another carbon adjacent to the point of attachment of Q^s is optionally substituted by R^{18} , a carbon adjacent to Q^b is optionally substituted by R^{16} , and another carbon adjacent to Q^b is optionally substituted by R^{19} ;

(ii) Y^{AT} wherein Y^{AT} is Q^b-Q^s ;

(iii) Q^b-Q^{ss} wherein Q^{ss} is $(CH(R^{14}))_e-W^2-(CH(R^{15}))_h$, wherein e and h are independently 1 or 2 and W^2 is $CR^{4a}=CR^{4b}$ with the proviso that $(CH(R^{14}))_e$ is bonded to E^0 ; and

(iv) Q^b-Q^{ssss} or Q^b-Q^{sssr} wherein Q^{ssss} is $(CH(R^{38}))_r-W^5$ and Q^{sssr} is $(CH(R^{38}))_r-W^6$, r is an integer selected from 1 through 2, W^5 and W^6 are independently selected from the group consisting of 1,4-indenyl, 1,5-indenyl, 1,6-indenyl, 1,7-indenyl, 2,7-indenyl, 2,6-indenyl, 2,5-indenyl, 2,4-indenyl, 3,4-indenyl, 3,5-indenyl, 3,6-indenyl, 3,7-indenyl, 2,4-benzofuranyl, 2,5-benzofuranyl, 2,6-benzofuranyl, 2,7-benzofuranyl, 3,4-benzofuranyl, 3,5-benzofuranyl, 3,6-

benzofuranyl, 3,7-benzofuranyl, 2,4-benzothiophenyl, 2,5-benzothiophenyl, 2,6-benzothiophenyl, 2,7-benzothiophenyl, 3,4-benzothiophenyl, 3,5-benzothiophenyl, 3,6-benzothiophenyl, 3,7-benzothiophenyl, 2,7-imidazo(1,2-a)pyridinyl, 3,4-imidazo(1,2-a)pyridinyl, 3,5-imidazo(1,2-a)pyridinyl, 3,6-imidazo(1,2-a)pyridinyl, 3,7-imidazo(1,2-a)pyridinyl, 2,4-indolyl, 2,5-indolyl, 2,6-indolyl, 2,7-indolyl, 3,4-indolyl, 3,5-indolyl, 3,6-indolyl, 3,7-indolyl, 1,4-isoindolyl, 1,5-isoindolyl, 1,6-isoindolyl, 2,4-isoindolyl, 2,5-isoindolyl, 2,6-isoindolyl, 2,7-isoindolyl, 1,3-isoindolyl, 3,4-indazolyl, 3,5-indazolyl, 3,6-indazolyl, 3,7-indazolyl, 2,4-benzoxazolyl, 2,5-benzoxazolyl, 2,6-benzoxazolyl, 2,7-benzoxazolyl, 3,4-benzisoxazolyl, 3,5-benzisoxazolyl, 3,6-benzisoxazolyl, 3,7-benzisoxazolyl, 1,4-naphthyl, 1,5-naphthyl, 1,6-naphthyl, 1,7-naphthyl, 1,8-naphthyl, 2,4-naphthyl, 2,5-naphthyl, 2,6-naphthyl, 2,7-naphthyl, 2,8-naphthyl, 2,4-quinolinyl, 2,5-quinolinyl, 2,6-quinolinyl, 2,7-quinolinyl, 2,8-quinolinyl, 3,4-quinolinyl, 3,5-quinolinyl, 3,6-quinolinyl, 3,7-quinolinyl, 3,8-quinolinyl, 4,5-quinolinyl, 4,6-quinolinyl, 4,7-quinolinyl, 4,8-quinolinyl, 1,4-isoquinolinyl, 1,5-isoquinolinyl, 1,6-isoquinolinyl, 1,7-isoquinolinyl, 1,8-isoquinolinyl, 3,4-isoquinolinyl, 3,5-isoquinolinyl, 3,6-isoquinolinyl, 3,7-isoquinolinyl, 3,8-isoquinolinyl, 4,5-isoquinolinyl, 4,6-isoquinolinyl, 4,7-isoquinolinyl, 4,8-isoquinolinyl, 3,4-cinnolinyl, 3,5-cinnolinyl, 3,6-cinnolinyl, 3,7-cinnolinyl, 3,8-cinnolinyl, 4,5-cinnolinyl, 4,6-cinnolinyl, 4,7-cinnolinyl, and 4,8-cinnolinyl, and each carbon and hydrido containing nitrogen member of the ring of the W^5 and of the ring of the W^6 , other than the points of attachment of W^5 and W^6 , is optionally substituted with one or more of the group consisting of R^9 , R^{10} , R^{11} , and R^{12} , with the proviso that Q^b is bonded to lowest number substituent position of each W^5 , with further proviso that Q^b is bonded to highest number substituent position of each W^6 , and with the additional proviso that $(CH(R^{38}))_r$ is bonded to E^0 ;

R^{16} , R^{17} , R^{18} , and R^{19} are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, nitro, alkoxyamino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, alkenyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, haloalkoxyalkyl, carboalkoxy, and cyano;

~~R^{18} or R^{19} is optionally selected from the group consisting of $NR^{20}R^{21}$, $N(R^{20})C(NR^{25})N(R^{23})(R^{24})$, and $C(NR^{25})NR^{23}R^{24}$, with the proviso that R^{18} , R^{19} , and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of $NR^{20}R^{21}$, aminoalkyl, hydrido, $N(R^{26})C(NR^{25})N(R^{23})(R^{24})$, and $C(NR^{25})NR^{23}R^{24}$, with the proviso that no more than one of R^{20} and R^{21} is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino ~~at the same time~~ and with the further proviso that no more than one of R^{23} and R^{24} is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino ~~at the same time~~;

R^{20} , R^{21} , R^{23} , R^{24} , R^{25} , and R^{26} are independently selected from the group consisting of hydrido, alkyl, hydroxy, aminoalkyl, amino, dialkylamino, alkylamino, and hydroxyalkyl;

Q^s is selected from the group consisting of a bond, $(CR^{37}R^{38})_b$ wherein b is an integer selected from 1 through 4, and $(CH(R^{14}))_c-W^1-(CH(R^{15}))_d$ wherein c and d are integers independently selected from 1 through 3 and W^1 is selected from the group consisting of $C(O)N(R^{14})$, $(R^{14})NC(O)$, $S(O)$, $S(O)_2$, $S(O)_2N(R^{14})$, $N(R^{14})S(O)_2$, and $N(R^{14})$, with the provisos that R^{14} is selected from other than halo when directly bonded to N and that $(CR^{37}R^{38})_b$ and $(CH(R^{14}))_c$ are bonded to E^0 ;

R^{14} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

R^{37} and R^{38} are independently selected from the group consisting of hydrido, alkyl, and haloalkyl; **and**

R^{38} is optionally aroyl or heteroaroyl, wherein R^{38} is optionally substituted with one or more substituents selected from the group consisting of R^{16} , R^{17} , R^{18} , and $R^{19}[[\cdot]]$.

Y^0 is optionally Y^{A^T} wherein Y^{A^T} is Q^b-Q^s ;


Y^0 is optionally Q^b-Q^{ss} wherein Q^{ss} is $(CH(R^{14}))_e-W^2-(CH(R^{15}))_h$ wherein e and h are independently 1 or 2 and W^2 is $CR^{4a}=CR^{4b}$ with the proviso that $(CH(R^{14}))_e$ is bonded to E^0 ;

Y^0 is optionally Q^b-Q^{ssss} or Q^b-Q^{sssr} wherein Q^{ssss} is

$(CH(R^{38}))_r-W^5$ and Q^{sssr} is $(CH(R^{38}))_r-W^6$, r is an integer selected from 1 through 2, W^5 and W^6 are independently selected from the group consisting of 1,4-indenyl, 1,5-indenyl, 1,6-indenyl, 1,7-indenyl, 2,7-indenyl, 2,6-indenyl, 2,5-indenyl, 2,4-indenyl, 3,4-indenyl, 3,5-indenyl, 3,6-indenyl, 3,7-indenyl, 2,4-benzofuranyl, 2,5-benzofuranyl, 2,6-benzofuranyl, 2,7-benzofuranyl, 3,4-benzofuranyl, 3,5-benzofuranyl, 3,6-benzofuranyl, 3,7-benzofuranyl, 2,4-benzothiophenyl, 2,5-benzothiophenyl, 2,6-benzothiophenyl, 2,7-benzothiophenyl, 3,4-benzothiophenyl, 3,5-benzothiophenyl, 3,6-benzothiophenyl, 3,7-benzothiophenyl, 2,7-imidazo(1,2-a)pyridinyl, 3,4-imidazo(1,2-a)pyridinyl, 3,5-imidazo(1,2-a)pyridinyl, 3,6-imidazo(1,2-a)pyridinyl, 3,7-imidazo(1,2-a)pyridinyl, 2,4-

~~indolyl, 2,5-indolyl, 2,6-indolyl, 2,7-indolyl, 3,4-indolyl, 3,5-indolyl, 3,6-indolyl, 3,7-indolyl, 1,4-isoindolyl, 1,5-isoindolyl, 1,6-isoindolyl, 2,4-isoindolyl, 2,5-isoindolyl, 2,6-isoindolyl, 2,7-isoindolyl, 1,3-isoindolyl, 3,4-indazolyl, 3,5-indazolyl, 3,6-indazolyl, 3,7-indazolyl, 2,4-benzoxazolyl, 2,5-benzoxazolyl, 2,6-benzoxazolyl, 2,7-benzoxazolyl, 3,4-benzisoxazolyl, 3,5-benzisoxazolyl, 3,6-benzisoxazolyl, 3,7-benzisoxazolyl, 1,4-naphthyl, 1,5-naphthyl, 1,6-naphthyl, 1,7-naphthyl, 1,8-naphthyl, 2,4-naphthyl, 2,5-naphthyl, 2,6-naphthyl, 2,7-naphthyl, 2,8-naphthyl, 2,4-quinoliny, 2,5-quinoliny, 2,6-quinoliny, 2,7-quinoliny, 2,8-quinoliny, 3,4-quinoliny, 3,5-quinoliny, 3,6-quinoliny, 3,7-quinoliny, 3,8-quinoliny, 4,5-quinoliny, 4,6-quinoliny, 4,7-quinoliny, 4,8-quinoliny, 1,4-isoquinoliny, 1,5-isoquinoliny, 1,6-isoquinoliny, 1,7-isoquinoliny, 1,8-isoquinoliny, 3,4-isoquinoliny, 3,5-isoquinoliny, 3,6-isoquinoliny, 3,7-isoquinoliny, 3,8-isoquinoliny, 4,5-isoquinoliny, 4,6-isoquinoliny, 4,7-isoquinoliny, 4,8-isoquinoliny, 3,4-cinnoliny, 3,5-cinnoliny, 3,6-cinnoliny, 3,7-cinnoliny, 3,8-cinnoliny, 4,5-cinnoliny, 4,6-cinnoliny, 4,7-cinnoliny, and 4,8-cinnoliny, and each carbon and hyrido-containing nitrogen member of the ring of the W^5 and of the ring of the W^6 , other than the points of attachment of W^5 and W^6 , is optionally substituted with one or more of the group consisting of R^9 , R^{10} , R^{11} , and R^{12} , with the proviso that Q^b is bonded to lowest number substituent position of each W^5 , with further proviso that Q^b is bonded to highest number substituent position of each W^6 , and with the additional proviso that $(CH(R^{38}))_r$ is bonded to E^6 .~~

2. (currently amended): The compound as recited in **[[Claim]] claim 1** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of:

(i) [[B is]] phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{32} , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally

substituted by R³⁶, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³², is optionally substituted by R³³, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³⁶, is optionally substituted by R³⁵, and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R³³ and R³⁵, respectively, is optionally substituted by R³⁴; a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³², the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R³⁶, a carbon adjacent to R³² and two atoms from the carbon at the point of attachment is optionally substituted by R³³, a carbon adjacent to R³⁶ and two atoms from the carbon at the point of attachment is optionally substituted by R³⁵, and any carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;

(ii) hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶; and

(iii) a C3-C12 cycloalkyl or C4-C9 heterocyclyl, wherein (a) each ring carbon may be optionally substituted with R₃₃, (b) a ring carbon, other than the ring carbon at the point of attachment of B to A, may be optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, (c) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (d) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (e) a ring carbon or nitrogen atom, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (f) a ring carbon or nitrogen, if present, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², (g) a ring carbon or nitrogen, if present, in a first gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally

substituted by R¹⁰, is optionally substituted by R¹¹, (h) a ring carbon or nitrogen, if present, in a second gamma position relative to the carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹², is optionally substituted by R³³, and (i) a ring carbon or nitrogen, if present, in a delta position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹¹ and R³³, respectively, is optionally substituted by R³⁴.

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkylenedioxy, haloalkylthio, alkanoyloxy, alkoxy, hydroxy, amino, alkoxyamino, haloalkanoyl, nitro, alkylamino, alkylthio, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkylsulfonamido, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, alkyl, alkenyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b;

~~B is optionally selected from the group consisting of hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;~~

~~B is optionally a C3-C12 cycloalkyl or C4-C9 heterocyclyl, wherein each ring carbon may be optionally substituted with R³³, a ring carbon other than the ring carbon at the point of attachment of B to A may be optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, ring carbons and nitrogens adjacent to the carbon at the point of attachment may be optionally substituted with R⁹ or R¹³, a ring carbon or nitrogen adjacent to the R⁹ position and two atoms from the point of attachment may be substituted with R¹⁰, a ring carbon or nitrogen adjacent to the R¹³ position and two atoms from the point of attachment may be substituted with R¹², a ring carbon three atoms from the point of attachment and adjacent to the R¹⁰ position may be substituted with R¹¹, a ring carbon three atoms from the point of attachment and adjacent to the R¹² position may be substituted with R³³, and a ring carbon four atoms from the point of attachment and adjacent to the R¹¹ and R³³ positions may be substituted with R³⁴;~~

R⁹, R¹⁰, R¹¹, R¹², and R¹³ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, alkoxyamino, alkanoyl, haloalkanoyl, amidino, guanidino, alkylenedioxy, haloalkylthio, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy,

aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkylamino, N-alkyl-N-arylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylthio, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, alkylsulfamido, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, amidosulfonyl, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, and cyano;

A is a bond or $(\text{CH}(\text{R}^{15}))_{\text{pa}}-(\text{W}^7)_{\text{r}}$ wherein r is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is selected from the group consisting of O, S, $\text{C}(\text{O})$, $(\text{R}^7)\text{NC}(\text{O})$, $(\text{R}^7)\text{NC}(\text{S})$, and $\text{N}(\text{R}^7)$;

R^7 is selected from the group consisting of hydrido, hydroxy and alkyl;

R^{15} is selected from the group consisting of hydrido, hydroxy, halo, alkyl, and haloalkyl;

M is N or $\text{R}^1\text{-C}$;

R^1 is selected from the group consisting of hydrido, alkyl, cyano, halo, haloalkyl, haloalkoxy, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

R^2 is $\text{Z}^0\text{-Q}$;

Z^0 is selected from the group consisting of:

(i) a bond, $\text{W}^0\text{-(CH}(\text{R}^{42}))_{\text{p}}$ wherein p is an integer selected from 0 through 3 and W^0 is selected from the group consisting of O, S, and $\text{N}(\text{R}^{41})$, and $(\text{CH}(\text{R}^{41}))_{\text{g}}\text{-O}$ wherein g is an integer selected from 1 through 3, with the proviso that Z^0 is directly bonded to the pyrimidinone ring; **and**

(ii) ~~Z^0 is optionally~~ $\text{W}^{22}\text{-(CH}(\text{R}^{42}))_{\text{h}}$ wherein h is 0 or 1 and W^{22} is selected from the group consisting of 1,2-cyclopropyl, 1,2-cyclobutyl, 1,2-cyclohexyl, 1,3-cyclohexyl, 1,2-cyclopentyl, 1,3-cyclopentyl, 2,3-morpholinyl, 2,4-morpholinyl, 2,6-morpholinyl, 3,4-morpholinyl, 3,5-morpholinyl, 1,2-piperazinyl, 1,3-piperazinyl, 2,3-piperazinyl, 2,6-piperazinyl, 1,2-piperidinyl, 1,3-piperidinyl, 2,3-piperidinyl, 2,4-piperidinyl, 2,6-piperidinyl, 3,4-piperidinyl, 1,2-pyrrolidinyl, 1,3-pyrrolidinyl, 2,3-pyrrolidinyl, 2,4-pyrrolidinyl, 2,5-pyrrolidinyl, 3,4-pyrrolidinyl, 2,3-tetrahydrofuranyl, 2,4-tetrahydrofuranyl, 2,5-tetrahydrofuranyl, and

3,4-tetrahydrofuranyl, wherein Z^0 is directly bonded to the pyrimidinone ring and W^{22} is optionally substituted with one or more substituents selected from the group consisting of R^9 , R^{10} , R^{11} , R^{12} , and R^{13} ;

R^{41} is selected from the group consisting of hydrido, hydroxy, amino, and alkyl;

R^{42} is selected from the group consisting of amidino, hydroxyamino, hydrido, hydroxy, amino, and alkyl;

Q is selected from the group consisting of:

(i) ~~[[Q is]] phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11} a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , the other carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^{13} , a carbon adjacent to R^9 and two atoms from the carbon at the point of attachment is optionally substituted by R^{10} , a carbon adjacent to R^{13} and two atoms from the carbon at the point of attachment is optionally substituted by R^{12} , and any carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than a phenyl when Z^0 is a bond; **and**~~


~~(ii) Q is optionally hydrido with the proviso that Z^0 is selected from other than a bond;~~

K is CHR^{4a} wherein R^{4a} is selected from the group consisting of hydrido, hydroxyalkyl, alkyl, alkoxyalkyl, alkylthioalkyl, and haloalkyl;

E^0 is selected from the group consisting of a bond, $C(O)N(H)$, $(H)NC(O)$, $(R^7)NS(O)_2$, and $S(O)_2N(R^7)$;

Y^0 is selected from the group consisting of:

(i) [[Y⁰ is]] the formula



wherein J⁵, J⁶, D⁵, D⁶ and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J⁵ and J⁶ is absent when T is a 5-membered heteroaryl ring, J⁵ is optionally substituted by R¹⁷ when J⁵ is a carbon atom, J⁶ is optionally substituted by R¹⁸ when J⁶ is a carbon atom, D⁵ is optionally substituted by R¹⁶ when D⁵ is a carbon atom and D⁶ is optionally substituted by R¹⁹ when D⁶ is a carbon atom; phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s, a carbon two or three contiguous atoms from the point of attachment of Q^s to the phenyl or heteroaryl ring is substituted by Q^b, a carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁷, another carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁸, a carbon adjacent to Q^b is optionally substituted by R¹⁶, and another carbon adjacent to Q^b is optionally substituted by R¹⁹;

(ii) Y^{AT} wherein Y^{AT} is Q^b-Q^s; and

(iii) Q^b-Q^{ss} wherein Q^{ss} is (CH(R¹⁴))_e-W²-(CH(R¹⁵))_h, wherein e and h are independently 1 or 2 and W² is CR^{4a}=CH with the proviso that (CH(R¹⁴))_e is bonded to E⁰;

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;
R¹⁶ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that no more than one of R²⁰ and R²¹ is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino at the same time and with the further proviso that no more than one of R²³

and R^{24} is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino ~~at the same time~~;

R^{20} , R^{21} , R^{23} , R^{24} , R^{25} , and R^{26} are independently selected from the group consisting of hydrido, alkyl, hydroxy, amino, alkylamino and dialkylamino;

Q^s is selected from the group consisting of a bond, $(CR^{37}R^{38})_b$ wherein b is an integer selected from 1 through 4, and $(CH(R^{14}))_c-W^1-(CH(R^{15}))_d$ wherein c and d are integers independently selected from 1 through 3 and W^1 is selected from the group consisting of $C(O)N(R^{14})$, $(R^{14})NC(O)$, $S(O)$, $S(O)_2$, $S(O)_2N(R^{14})$, $N(R^{14})S(O)_2$, and $N(R^{14})$, with the proviso that R^{14} is selected from other than halo when directly bonded to N and with the further proviso that $(CR^{37}R^{38})_b$ and $(CH(R^{14}))_c$ are bonded to E^0 ;

R^{14} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

R^{37} and R^{38} are independently selected from the group consisting of hydrido, alkyl, and haloalkyl; **and**

R^{38} is optionally aroyl or heteroaroyl, wherein R^{38} is optionally substituted with one or more substituents selected from the group consisting of R^{16} , R^{17} , R^{18} , and R^{19} [:];

~~Y^0 is optionally Y^{AT} wherein Y^{AT} is Q^b-Q^s ;~~

~~Y^0 is optionally Q^b-Q^{ss} wherein Q^{ss} is $(CH(R^{14}))_e-W^2-(CH(R^{15}))_h$, wherein e and h are independently 1 or 2 and W^2 is $CR^{4a}=CH$ with the proviso that $(CH(R^{14}))_e$ is bonded to E^0 .~~

3. (currently amended): The compound as recited in **[[Claim]] claim 2** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} ;

R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b ;

A is $(\text{CH}(\text{R}^{15}))_{\text{pa}}-\text{W}^7$ wherein pa is an integer selected from 0 through 3 and W^7 is selected from the group consisting of O, S, and $\text{N}(\text{R}^7)$ wherein R^7 is hydrido or alkyl;

R^{15} is selected from the group consisting of hydrido, hydroxy, halo, alkyl, and haloalkyl with the proviso that R^{15} is other than hydroxy and halo when R^{15} is on the carbon bonded directly to W^7 ;

M is N or R^1-C ;

R^1 is selected from the group consisting of hydrido, alkyl, cyano, halo, haloalkyl, haloalkoxy, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

R^2 is Z^0-Q ;

Z^0 is a bond or $\text{W}^0-(\text{CH}(\text{R}^{42}))_p$ wherein p is an integer selected from 0 through 3 and W^0 is selected from the group consisting of O, S, and $\text{N}(\text{R}^{41})$, with the proviso that Z^0 is directly bonded to the pyrimidinone ring;

R^{41} is selected from the group consisting of hydrido, hydroxy, and alkyl;

R^{42} is selected from the group consisting of amidino, hydrido, hydroxy, amino, and alkyl;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11}** a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R^{13} , a carbon adjacent to R^9 and two atoms from the carbon at the point of attachment is optionally substituted by R^{10} , a carbon adjacent to R^{13} and two atoms from the carbon at the point of attachment is optionally

substituted by R^{12} , and any carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than a phenyl when Z^0 is a bond;


R^9 , R^{10} , R^{11} , R^{12} , and R^{13} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, alkoxyamino, alkanoyl, haloalkanoyl, amidino, guanidino, alkylenedioxy, haloalkylthio, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocycloxy, heterocyclylalkoxy, hydroxy, amino, alkylamino, N-alkyl-N-aryl amino, arylamino, aralkylamino, heteroaryl amino, heteroaralkyl amino, heterocyclyl amino, heterocyclylalkyl amino, alkylthio, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, alkylsulfamido, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, amidosulfonyl, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, and cyano;

K is CHR^{4a} wherein R^{4a} is selected from the group consisting of hydrido, hydroxyalkyl, alkyl, alkoxyalkyl, alkylthioalkyl, and haloalkyl;

E^0 is selected from the group consisting of a bond, $C(O)N(H)$, $(H)NC(O)$, $(R^7)NS(O)_2$, and $S(O)_2N(R^7)$;

Y^0 is selected from the group consisting of:

(i) $[[Y^0 \text{ is}]]$ the formula



wherein J^5 , J^6 , D^5 , D^6 and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J^5 and J^6 is absent when T is a 5-membered heteroaryl ring, J^5 is optionally substituted by R^{17} when J^5 is a carbon atom, J^6 is optionally substituted by R^{18} when J^6 is a carbon atom, D^5 is optionally substituted by R^{16} when D^5 is a carbon atom and D^6 is optionally substituted by R^{19} when D^6 is a carbon atom; phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^a , a carbon two or three contiguous atoms from the point of attachment of Q^a to the phenyl or heteroaryl ring is substituted by Q^b , a

~~carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁷;
another carbon adjacent to the point of attachment of Q^s is optionally substituted by
R¹⁸; a carbon adjacent to Q^b is optionally substituted by R¹⁶; and another carbon
adjacent to Q^b is optionally substituted by R¹⁹; and~~

**(ii) Q^b-Q^{ss} wherein Q^{ss} is (CH(R¹⁴))_e-W²-(CH(R¹⁵))_h, wherein e and h
are integers independently selected from 1 through 2 and W² is CR^{4a}=CH with
the proviso that (CH(R¹⁴))_e is bonded to E⁰;**

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of
hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino,
alkoxyamino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl,
alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;

~~R¹⁶ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹,
N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b
are not simultaneously hydrido;—~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido,
N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that no more than one
of R²⁰ and R²¹ is selected from the group consisting of hydroxy, amino, alkylamino, and
dialkylamino ~~at the same time~~ and with the further proviso that no more than one of R²³
and R²⁴ is selected from the group consisting of hydroxy, amino, alkylamino, and
dialkylamino ~~at the same time~~;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group
consisting of hydrido, alkyl, hydroxy, amino, alkylamino and dialkylamino;

Q^s is selected from the group consisting of a bond, (CR³⁷R³⁸)_b, wherein b is an
integer selected from 1 through 3, and (CH(R¹⁴))_c-W¹-(CH(R¹⁵))_d, wherein c and d
are independently 1 or 2 and W¹ is selected from the group consisting of
C(O)N(R¹⁴), (R¹⁴)NC(O), S(O), S(O)₂, S(O)₂N(R¹⁴), N(R¹⁴)S(O)₂, and N(R¹⁴), with the
proviso that R¹⁴ is selected from other than halo when directly bonded to N and with
the further proviso that (CR³⁷R³⁸)_b, and (CH(R¹⁴))_c are bonded to E⁰;

R¹⁴ is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;


R¹⁴ is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

R³⁷ and R³⁸ are independently selected from the group consisting of hydrido,
alkyl, and haloalkyl; **and**

R³⁸ is optionally aroyl and heteroaroyl[;].

~~Y⁰ is optionally Q^b-Q^{ss} wherein Q^{ss} is (CH(R¹⁴))_e-W²-(CH(R¹⁵))_h, wherein e and h are integers independently selected from 1 through 2 and W² is CR⁴²=CH with the proviso that (CH(R¹⁴))_e is bonded to E⁰.~~

4. (currently amended): The compound as recited in **[[Claim]] claim 3** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, trialkylsilyl, C2-C4 alkyl, C3-C5 alkylenyl, C3-C4 alkenyl, C3-C4 alkynyl, and C2-C4 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 3 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, and R³⁴;

R³², R³³, and R³⁴ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboalkoxy, carboxy, carboxamido, and cyano;

A is (CH(R¹⁵))_{pa}-N(R⁷) wherein pa is an integer selected from 0 through 2 and R⁷ is selected from the group consisting of hydrido and alkyl;

R¹⁵ is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, alkyl, cyano, halo, haloalkyl, haloalkoxy, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

R² is Z⁰-Q;

Z⁰ is a bond or W⁰-CH(R⁴²) wherein W⁰ is selected from the group consisting of O, S, and N(R⁴¹);


R⁴¹ and R⁴² are independently hydrido or alkyl;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹ ~~a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond;~~

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkylsulfonamido, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, alkyl, alkoxy, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkoxyamino, alkylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylsulfonamido, amidosulfonyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, halo, haloalkyl, and cyano;

Y⁰ is the formula



wherein J⁵, J⁶, D⁵, D⁶ and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J⁵ and J⁶ is absent when T is a 5-membered heteroaryl ring, J⁵ is optionally substituted by R¹⁷ when J⁵ is a carbon atom, J⁶ is optionally substituted by R¹⁸ when J⁶ is a carbon atom, D⁵ is optionally substituted by R¹⁶ when D⁵ is a carbon atom and D⁶ is optionally substituted by R¹⁹ when D⁶ is a carbon atom;
~~phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s, a carbon two or three contiguous atoms from the point of attachment of Q^s to the phenyl or heteroaryl ring is substituted by Q^b, a carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁷, another carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁸, a carbon adjacent to Q^b is optionally substituted by R¹⁸, and another carbon adjacent to Q^b is optionally substituted by R¹⁹;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;

~~R¹⁸ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁸, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that no more than one of R²⁰ and R²¹ is hydroxy ~~at the same time~~ and with the further proviso that no more than one of R²³ and R²⁴ is hydroxy ~~at the same time~~;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, alkyl, and hydroxy; **and**

Q^s is selected from the group consisting of abond, CH₂, and CH₂CH₂.

5. (currently amended): Compound of **[[Claim]] claim 4** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, butyl, 2-butenyl, 3-butenyl, 2-butylnyl, sec-butyl, tert-butyl, isobutyl, 2-methylpropenyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, and 2,2-difluoropropyl, wherein each member of group B is optionally substituted at any carbon up to and including 3 atoms from the point of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , and R^{34} ;

R^{32} , R^{33} , and R^{34} are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, amidosulfonyl, N-methylamidossulfonyl, N,N-dimethylamidossulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;

A is selected from the group consisting of a bond, NH, and $\text{N}(\text{CH}_3)$;

M is N or $\text{R}^1\text{-C}$;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, 1-aminoethyl, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, methoxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxyamino, methylthio, ethylthio, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, and bromo;

R^2 is $\text{Z}^0\text{-Q}$;

Z^0 is selected from the group consisting of a bond, O, S, NH, $\text{N}(\text{CH}_3)$, OCH_2 , SCH_2 , $\text{N}(\text{H})\text{CH}_2$, and $\text{N}(\text{CH}_3)\text{CH}_2$;

Q is selected from the group consisting of phenyl **[[.]] and** 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, and 1,3,5-triazin-2-yl **heteroaryl rings**, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is**

optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹ a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond;




R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;





R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, carboxymethyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl,

N,N-dimethylamidodisulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl, N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidodisulfonyl, N-(2-chlorobenzyl)amidodisulfonyl, N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl, N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl, N-cyclohexylamidocarbonyl, fluoro, chloro, bromo, cyano, cyclobutoxy, cyclohexoxy, cyclohexylmethoxy, 4-trifluoromethylcyclohexylmethoxy, cyclopentoxymethyl, benzyl, benzyloxy, 4-bromo-3-fluorophenoxy, 3-bromobenzyloxy, 4-bromobenzyloxy, 4-bromobenzylamino, 5-bromopyrid-2-ylmethylamino, 4-butoxyphenylamino, 3-chlorobenzyl, 4-chlorophenoxy, 4-chloro-3-ethylphenoxy, 4-chloro-3-ethylbenzylamino, 4-chloro-3-ethylphenylamino, 3-chlorobenzyloxy, 4-chlorobenzyloxy, 4-chlorobenzylsulfonyl, 4-chlorophenylamino, 4-chlorophenylsulfonyl, 5-chloropyrid-3-yl, 2-cyanopyrid-3-yl, 2,3-difluorobenzyloxy, 2,4-difluorobenzyloxy, 3,4-difluorobenzyloxy, 2,5-difluorobenzyloxy, 3,5-difluorophenoxy, 3,5-difluorobenzyloxy, 4-difluoromethoxybenzyloxy, 2,3-difluorophenoxy, 2,4-difluorophenoxy, 2,5-difluorophenoxy, 3,5-dimethylphenoxy, 3,4-dimethylphenoxy, 3,4-dimethylbenzyloxy, 3,5-dimethylbenzyloxy, 4-ethoxyphenoxy, 4-ethylbenzyloxy, 3-ethylphenoxy, 4-ethylaminophenoxy, 3-ethyl-5-methylphenoxy, 4-fluorobenzyloxy, 2-fluoro-3-trifluoromethylbenzyloxy, 3-fluoro-5-trifluoromethylbenzyloxy, 4-fluoro-2-trifluoromethylbenzyloxy, 4-fluoro-3-trifluoromethylbenzyloxy, 2-fluorophenoxy, 4-fluorophenoxy, 2-fluoro-3-trifluoromethylphenoxy, 2-fluorobenzyloxy, 4-fluorophenylamino, 2-fluoro-4-trifluoromethylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, phenylamino, 1-phenylethoxy, 2-phenylethoxy, 2-phenylethyl, 2-phenylethylamino, phenylsulfonyl, 3-trifluoromethoxybenzyloxy, 4-trifluoromethoxybenzyloxy, 3-trifluoromethoxyphenoxy, 4-trifluoromethoxyphenoxy,

3-trifluoromethylbenzyloxy, 4-trifluoromethylbenzyloxy,
2,4-bis-trifluoromethylbenzyloxy, 3-trifluoromethylbenzyl,
3,5-bis-trifluoromethylbenzyloxy, 4-trifluoromethylphenoxy,
3-trifluoromethylphenoxy, 3-trifluoromethylthiobenzyloxy,
4-trifluoromethylthiobenzyloxy, 2,3,4-trifluorophenoxy, 2,3,5-trifluorophenoxy, 3-
pentafluoroethylphenoxy, 3-(1,1,2,2-tetrafluoroethoxy)phenoxy, and
3-trifluoromethylthiophenoxy;

Y^0 is selected from the group consisting of:





~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine,
3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine, 2-Q^b-5-Q^s-3-R¹⁶-6-R¹⁸pyrazine, 3-Q^b-6-Q^s-2-R¹⁶-
5-R¹⁸-4-R¹⁹pyridazine, 2-Q^b-5-Q^s-4-R¹⁷-6-R¹⁸pyrimidine, 5-Q^b-2-Q^s-4-R¹⁶-6-
R¹⁹pyrimidine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹thiophene, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷thiophene,
3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹furan, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷furan, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹pyrrole,
2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷pyrrole, 4-Q^b-2-Q^s-5-R¹⁹imidazole, 2-Q^b-4-Q^s-5-R¹⁷imidazole,
3-Q^b-5-Q^s-4-R¹⁶isoxazole, 5-Q^b-3-Q^s-4-R¹⁶isoxazole, 2-Q^b-5-Q^s-4-R¹⁶pyrazole, 4-Q^b-
2-Q^s-5-R¹⁹thiazole, and 2-Q^b-5-Q^s-4-R¹⁷thiazole;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, isopropyl, propyl, carboxy, amidino, guanidino, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl, ethylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, and cyano;


~~R¹⁶ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁰)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that no more than one of R²⁰, R²¹, R²³, and R²⁴ can be hydroxy, when any two of the group consisting of R²⁰, R²¹, R²³, and R²⁴ are bonded to the same atom and with the further proviso that said Q^b group is bonded directly to a carbon atom;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, methyl, ethyl, propyl, butyl, isopropyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH₂, and CH₂CH₂.

6. (currently amended): The compound as recited in **[[Claim]] claim 4** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

A is selected from the group consisting of $\text{CH}_2\text{N}(\text{CH}_3)$, $\text{CH}_2\text{N}(\text{CH}_2\text{CH}_3)$, $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)$, and $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_3)$;

M is N or $\text{R}^1\text{-C}$;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, 1-aminoethyl, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, methoxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxyamino, methylthio, ethylthio, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, and bromo;

R^2 is $\text{Z}^0\text{-Q}$;

Z^0 is selected from the group consisting of a bond, O, S, NH, $\text{N}(\text{CH}_3)$, OCH_2 , SCH_2 , $\text{N}(\text{H})\text{CH}_2$, and $\text{N}(\text{CH}_3)\text{CH}_2$;

Q is selected from the group consisting of phenyl **[[,]] and** 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, and 1,3,5-triazin-2-yl **heteroaryl rings**, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the**





point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹ a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond;




R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, methanesulfonamido, amidosulfonyl, N-methylamidossulfonyl, N,N-dimethylamidossulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, carboxymethyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methanesulfonamido, amidosulfonyl, N-methylamidossulfonyl, N,N-dimethylamidossulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl, N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidossulfonyl, N-(2-chlorobenzyl)amidossulfonyl, N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl,

N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl, N-cyclohexylamidocarbonyl, fluoro, chloro, bromo, cyano, cyclobutoxy, cyclohexoxy, cyclohexylmethoxy, 4-trifluoromethylcyclohexylmethoxy, cyclopentoxo, benzyl, benzyloxy, 4-bromo-3-fluorophenoxy, 3-bromobenzyloxy, 4-bromobenzyloxy, 4-bromobenzylamino, 5-bromopyrid-2-ylmethylamino, 4-butoxyphenamino, 3-chlorobenzyl, 4-chlorophenoxy, 4-chloro-3-ethylphenoxy, 4-chloro-3-ethylbenzylamino, 4-chloro-3-ethylphenylamino, 3-chlorobenzyloxy, 4-chlorobenzyloxy, 4-chlorobenzylsulfonyl, 4-chlorophenylamino, 4-chlorophenylsulfonyl, 5-chloropyrid-3-ylloxy, 2-cyanopyrid-3-ylloxy, 2,3-difluorobenzyloxy, 2,4-difluorobenzyloxy, 3,4-difluorobenzyloxy, 2,5-difluorobenzyloxy, 3,5-difluorophenoxy, 3,5-difluorobenzyloxy, 4-difluoromethoxybenzyloxy, 2,3-difluorophenoxy, 2,4-difluorophenoxy, 2,5-difluorophenoxy, 3,5-dimethylphenoxy, 3,4-dimethylphenoxy, 3,4-dimethylbenzyloxy, 3,5-dimethylbenzyloxy, 4-ethoxyphenoxy, 4-ethylbenzyloxy, 3-ethylphenoxy, 4-ethylaminophenoxy, 3-ethyl-5-methylphenoxy, 4-fluorobenzyloxy, 2-fluoro-3-trifluoromethylbenzyloxy, 3-fluoro-5-trifluoromethylbenzyloxy, 4-fluoro-2-trifluoromethylbenzyloxy, 4-fluoro-3-trifluoromethylbenzyloxy, 2-fluorophenoxy, 4-fluorophenoxy, 2-fluoro-3-trifluoromethylphenoxy, 2-fluorobenzyloxy, 4-fluorophenylamino, 2-fluoro-4-trifluoromethylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, phenylamino, 1-phenylethoxy, 2-phenylethoxy, 2-phenylethyl, 2-phenylethylamino, phenylsulfonyl, 3-trifluoromethoxybenzyloxy, 4-trifluoromethoxybenzyloxy, 3-trifluoromethoxyphenoxy, 4-trifluoromethoxyphenoxy, 3-trifluoromethylbenzyloxy, 4-trifluoromethylbenzyloxy, 2,4-bis-trifluoromethylbenzyloxy, 3-trifluoromethylbenzyl, 3,5-bis-trifluoromethylbenzyloxy, 4-trifluoromethylphenoxy, 3-trifluoromethylphenoxy, 3-trifluoromethylthiobenzyloxy, 4-trifluoromethylthiobenzyloxy, 2,3,4-trifluorophenoxy, 2,3,5-trifluorophenoxy, 3-pentafluoroethylphenoxy, 3-(1,1,2,2-tetrafluoroethoxy)phenoxy, and 3-trifluoromethylthiophenoxy;

Y⁰ is selected from the group consisting of:





~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹ benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹ pyridine, 3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹ pyridine, 2-Q^b-5-Q^s-3-R¹⁶-6-R¹⁸ pyrazine, 3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹ pyridazine, 2-Q^b-5-Q^s-4-R¹⁷-6-R¹⁸ pyrimidine, 5-Q^b-2-Q^s-4-R¹⁶-6-R¹⁹ pyrimidine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹ thiophene, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷ thiophene, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹ furan, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷ furan, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹ pyrrole, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷ pyrrole, 4-Q^b-2-Q^s-5-R¹⁹ imidazole, 2-Q^b-4-Q^s-5-R¹⁷ imidazole, 3-Q^b-5-Q^s-4-R¹⁶ isoxazole, 5-Q^b-3-Q^s-4-R¹⁶ isoxazole, 2-Q^b-5-Q^s-4-R¹⁶ pyrazole, 4-Q^b-2-Q^s-5-R¹⁹ thiazole, and 2-Q^b-5-Q^s-4-R¹⁷ thiazole;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, isopropyl, propyl, carboxy, amidino, guanidino, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio,

isopropylthio, trifluoromethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl, ethylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, and cyano;

Q^b is selected from the group consisting of $NR^{20}R^{21}$, $C(NR^{25})NR^{23}R^{24}$, and $N(R^{26})C(NR^{25})N(R^{23})(R^{24})$, with the proviso that no more than one of R^{20} , R^{21} , R^{23} , and R^{24} can be hydroxy, when any two of the group consisting of R^{20} , R^{21} , R^{23} , and R^{24} are bonded to the same atom and with the further proviso that said Q^b group is bonded directly to a carbon atom;

R^{20} , R^{21} , R^{23} , R^{24} , R^{25} , and R^{26} are independently selected from the group consisting of hydrido, methyl, ethyl, propyl, butyl, isopropyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH_2 , and CH_2CH_2 .

7. (currently amended): The compound as recited in **[[Claim]] claim 6** or a pharmaceutically acceptable salt thereof, wherein;

A is selected from the group consisting of $CH_2N(CH_3)$, $CH_2N(CH_2CH_3)$, $CH_2CH_2N(CH_3)$, and $CH_2CH_2N(CH_2CH_3)$;

M is N or R^1-C ;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy, hydroxymethyl, methoxyamino, methylthio, trifluoromethoxy, fluoro, and chloro;

R^2 is Z^0-Q ;


Z^0 is selected from the group consisting of a bond, O, S, NH, and $N(CH_3)$;


Q is selected from the group consisting of

3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-benzylphenyl, 3-amino-5-(2-phenylethyl)phenyl, 3-amino-5-benzylaminophenyl, 3-amino-5-(2-phenylethylamino)phenyl, 3-amino-5-benzyloxyphenyl, 3-amino-5-(2-phenylethoxy)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-benzylamidosulfonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl,

3-amino-5-(N-ethylamidocarbonyl)phenyl,
3-amino-5-(N-isopropylamidocarbonyl)phenyl,
3-amino-5-(N-propylamidocarbonyl)phenyl,
3-amino-5-(N-isobutylamidocarbonyl)phenyl,
3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl,
3-amino-5-(N-cyclobutylamidocarbonyl)phenyl,
3-amino-5-(N-cyclopentylamidocarbonyl)phenyl,
3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl,
3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl,
3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl,
-aminophenyl, 3-amino-5-(4-trifluoromethylbenzylamino)phenyl,
3-amino-5-(4-trifluoromethylbenzyloxy)phenyl, 3-carboxyphenyl,
3-carboxy-5-hydroxyphenyl, 3-amino-5-carboxyphenyl, 3-chlorophenyl,
2-chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl,
2-fluorophenyl, 3-fluorophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl,
3-methanesulfonylaminophenyl, 2-methoxyphenyl, 3-methoxyphenyl,
3-methoxyaminophenyl, 3-methoxycarbonylphenyl, 2-methylaminophenyl,
3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-
trifluoroacetamidophenyl, 3-trifluoromethylphenyl,
2-trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl,
3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl, with the proviso that
Q is other than a phenyl or a substituted phenyl when Z⁰ is a bond;

Y⁰ is selected from the group consisting of:





~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine,
3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹thiophene, and 2-Q^b-5-Q^s-
3-R¹⁸-4-R¹⁷thiophene;~~

R¹⁶ and R¹⁹ are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano;

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano;

Q^b is C(NR²⁵)NR²³R²⁴;

R²³, R²⁴, and R²⁵ are independently hydrido or methyl; **and**

Q^s is CH₂.

8. (currently amended): A compound as recited in **[[Claim]] claim 7** or a pharmaceutically acceptable salt thereof where said compound is selected from the group consisting of:

2-[3-[2-[3-aminophenoxy]-6-chloro-N-[[4-aminoiminomethylphenyl]methyl]-5-[N,N-dimethylhydrazino]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

2-[3-[2-[3-aminophenoxy]-6-chloro-5-[N-ethyl-N-methylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

2-[3-[2-[3-aminophenoxy]-6-chloro-5-[N,N-diethylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

[2-[4-[3-[3-aminophenoxy]-N-[[4-aminoiminomethylphenyl]methyl]-6-[N,N-dimethylhydrazino]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide;

2-[4-[3-[3-aminophenoxy]-6-[N-ethyl-N-methylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide;

2-[4-[3-[3-aminophenoxy]-6-[N,N-diethylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide;]

2-[3-[2-[3-amino-5-carboxyphenoxy]-6-chloro-N-[[4-aminoiminomethylphenyl]methyl]-5-[N,N-dimethylhydrazino]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

2-[3-[2-[3-amino-5-carboxyphenoxy]-6-chloro-5-[N-ethyl-N-methylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;


2-[3-[2-[3-amino-5-carboxyphenoxy]-6-chloro-5-[N,N-diethylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

[2-[4-[3-[3-amino-5-carboxyphenoxy]-N-[[4-aminoiminomethylphenyl]methyl]-6-[N,N-dimethylhydrazino]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide;

2-[4-[3-[3-amino-5-carboxyphenoxy]-6-[N-ethyl-N-methylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide; **and**

2-[4-[3-[3-amino-5-carboxyphenoxy]-6-[N,N-diethylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide.]

9. (currently amended): The compound as recited in **[[Claim]] claim 2** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is phenyl or a heteroaryl of 5 or 6 ring members, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³², (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³⁶, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³², is optionally substituted by R³³, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³⁶, is optionally substituted by R³⁵, and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha**

position relative to each of the ring atoms optionally substituted by R³³ and R³⁵, respectively, is optionally substituted by R³⁴; a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³²; the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R³⁶; a carbon adjacent to R³² and two atoms from the carbon at the point of attachment is optionally substituted by R³³; a carbon adjacent to R³⁶ and two atoms from the carbon at the point of attachment is optionally substituted by R³⁵; and any carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b;

A is a **bond** or (CH(R¹⁵))_{pa}-(W⁷)_{rr} wherein rr is 0 or 1, pa is an integer selected from 0 through 3, and W⁷ is (R⁷)NC(O) or N(R⁷);

R⁷ is selected from the group consisting of hydrido, hydroxy and alkyl;

R¹⁵ is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

R² is Z⁰-Q;

Z⁰ is a bond or W⁰-(CH(R⁴²))_p wherein p is 0 or 1 and W⁰ is selected from the group consisting of O, S, and N(R⁴¹);

R⁴¹ and R⁴² are independently hydrido or alkyl;


Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³,**

is optionally substituted by R¹², and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹ a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkylsulfonamido, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, alkyl, alkoxy, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkoxyamino, alkylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylsulfonamido, amidosulfonyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, halo, haloalkyl, and cyano;

Y⁰ is the formula



wherein J⁵, J⁶, D⁵, D⁶ and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J⁵ and J⁶ is absent when T is a 5-membered heteroaryl ring, J⁵ is optionally

substituted by R¹⁷ when J⁵ is a carbon atom, J⁶ is optionally substituted by R¹⁸ when J⁶ is a carbon atom, D⁵ is optionally substituted by R¹⁶ when D⁵ is a carbon atom and D⁶ is optionally substituted by R¹⁹ when D⁶ is a carbon atom;
~~phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s, a carbon two or three contiguous atoms from the point of attachment of Q^s to the phenyl or heteroaryl ring is substituted by Q^b, a carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁷, another carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁸, a carbon adjacent to Q^b is optionally substituted by R¹⁶, and another carbon adjacent to Q^b is optionally substituted by R¹⁹;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;

~~R¹⁶ or R¹⁹ is optionally NR²⁰R²¹ or C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydride;~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, and C(NR²⁵)NR²³R²⁴, with the proviso that no more than one of R²⁰ and R²¹ is hydroxy at the same time and with the further proviso that no more than one of R²³ and R²⁴ is hydroxy at the same time;

R²⁰, R²¹, R²³, R²⁴, and R²⁵ are independently selected from the group consisting of hydrido, alkyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH₂, and CH₂CH₂.

10. (currently amended): The compound as recited in **[[Claim]] claim 9** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of phenyl **and** **[[,]]** 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl **heteroaryl rings**, and 1,3,5-triazin-2-yl, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³², (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³⁶, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of**

attachment and in an alpha position relative to the ring atom optionally substituted by R³², is optionally substituted by R³³, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³⁶, is optionally substituted by R³⁵, and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R³³ and R³⁵, respectively, is optionally substituted by R³⁴; a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³², the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R³⁶, a carbon adjacent to R³² and two atoms from the carbon at the point of attachment is optionally substituted by R³³, a carbon adjacent to R³⁶ and two atoms from the carbon at the point of attachment is optionally substituted by R³⁵, and any carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, amidosulfonyl, N-methylamid sulfonyl, N,N-dimethylamid sulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, cyano, and Q^b;

A is selected from the group consisting of a bond, NH, N(CH₃), N(OH), CH₂, CH₃CH, CF₃CH, NHC(O), N(CH₃)C(O), C(O)NH, C(O)N(CH₃), CH₂CH₂, CH₂CH₂CH₂, CH₃CHCH₂, and CF₃CHCH₂;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, 1-aminoethyl, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, methoxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxyamino, methylthio, ethylthio, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, and bromo;

R^2 is Z^0 -Q;

Z^0 is selected from the group consisting of a bond, O, S, NH, $N(CH_3)$, OCH_2 , SCH_2 , $N(H)CH_2$, and $N(CH_3)CH_2$;

Q is selected from the group consisting of phenyl **and** **[[,]]** 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, and 1,3,5-triazin-2-yl **heteroaryl rings**, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11}** a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R^{13} , a carbon adjacent to R^9 and two atoms from the carbon at the point of attachment is optionally substituted by R^{10} , a carbon adjacent to R^{13} and two atoms from the carbon at the point of attachment is optionally substituted by R^{12} , and any carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than a phenyl when Z^0 is a bond;



R^9 , R^{11} , and R^{13} are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, methanesulfonamido, amidosulfonyl, N-methylamidossulfonyl, N,N-





dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;


R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, carboxymethyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl, N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidosulfonyl, N-(2-chlorobenzyl)amidosulfonyl, N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl, N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl, N-cyclohexylamidocarbonyl, fluoro, chloro, bromo, cyano, cyclobutoxy, cyclohexoxy, cyclohexylmethoxy, 4-trifluoromethylcyclohexylmethoxy, cyclopentoxy, benzyl, benzyloxy, 4-bromo-3-fluorophenoxy, 3-bromobenzyloxy, 4-bromobenzyloxy, 4-bromobenzylamino, 5-bromopyrid-2-ylmethylamino, 4-butoxyphenamino, 3-chlorobenzyl, 4-chlorophenoxy, 4-chloro-3-ethylphenoxy, 4-chloro-3-ethylbenzylamino, 4-chloro-3-ethylphenylamino, 3-chlorobenzyloxy, 4-chlorobenzyloxy, 4-chlorobenzylsulfonyl, 4-chlorophenylamino, 4-chlorophenylsulfonyl, 5-chloropyrid-3-yloxy, 2-cyanopyrid-3-yloxy, 2,3-difluorobenzyloxy, 2,4-difluorobenzyloxy, 3,4-difluorobenzyloxy, 2,5-difluorobenzyloxy, 3,5-difluorophenoxy, 3,5-difluorobenzyloxy, 4-difluoromethoxybenzyloxy, 2,3-difluorophenoxy, 2,4-difluorophenoxy, 2,5-difluorophenoxy, 3,5-dimethylphenoxy, 3,4-dimethylphenoxy, 3,4-dimethylbenzyloxy, 3,5-dimethylbenzyloxy, 4-ethoxyphenoxy, 4-ethylbenzyloxy, 3-ethylphenoxy, 4-ethylaminophenoxy, 3-ethyl-5-methylphenoxy, 4-fluorobenzyloxy, 2-fluoro-3-trifluoromethylbenzyloxy, 3-fluoro-5-trifluoromethylbenzyloxy,

4-fluoro-2-trifluoromethylbenzyloxy, 4-fluoro-3-trifluoromethylbenzyloxy, 2-fluorophenoxy, 4-fluorophenoxy, 2-fluoro-3-trifluoromethylphenoxy, 2-fluorobenzyloxy, 4-fluorophenylamino, 2-fluoro-4-trifluoromethylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, phenylamino, 1-phenylethoxy, 2-phenylethoxy, 2-phenylethyl, 2-phenylethylamino, phenylsulfonyl, 3-trifluoromethoxybenzyloxy, 4-trifluoromethoxybenzyloxy, 3-trifluoromethoxyphenoxy, 4-trifluoromethoxyphenoxy, 3-trifluoromethylbenzyloxy, 4-trifluoromethylbenzyloxy, 2,4-bis-trifluoromethylbenzyloxy, 3-trifluoromethylbenzyl, 3,5-bis-trifluoromethylbenzyloxy, 4-trifluoromethylphenoxy, 3-trifluoromethylphenoxy, 3-trifluoromethylthiobenzyloxy, 4-trifluoromethylthiobenzyloxy, 2,3,4-trifluorophenoxy, 2,3,5-trifluorophenoxy, 3-pentafluoroethylphenoxy, 3-(1,1,2,2-tetrafluoroethoxy)phenoxy, and 3-trifluoromethylthiophenoxy;

Y⁰ is selected from the group consisting of:







~~1-Q^b-4-Q^s-2-R¹⁸-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine, 3-Q^b-6-Q^s-2-R¹⁸-5-R¹⁸-4-R¹⁹pyridine, 2-Q^b-5-Q^s-3-R¹⁸-6-R¹⁸pyrazine, 3-Q^b-6-Q^s-2-R¹⁸-5-R¹⁸-4-R¹⁹pyridazine, 2-Q^b-5-Q^s-4-R¹⁷-6-R¹⁸pyrimidine, 5-Q^b-2-Q^s-4-R¹⁸-6-R¹⁹pyrimidine, 3-Q^b-5-Q^s-4-R¹⁸-2-R¹⁹thiophene, 2-Q^b-5-Q^s-3-R¹⁸-4-R¹⁷thiophene, 3-Q^b-5-Q^s-4-R¹⁸-2-R¹⁹furan, 2-Q^b-5-Q^s-3-R¹⁸-4-R¹⁷furan, 3-Q^b-5-Q^s-4-R¹⁸-2-R¹⁹pyrrole, 2-Q^b-5-Q^s-3-R¹⁸-4-R¹⁷pyrrole, 4-Q^b-2-Q^s-5-R¹⁹imidazole, 2-Q^b-4-Q^s-5-R¹⁷imidazole, 3-Q^b-5-Q^s-4-R¹⁸isoxazole, 5-Q^b-3-Q^s-4-R¹⁸isoxazole, 2-Q^b-5-Q^s-4-R¹⁸pyrazole, 4-Q^b-2-Q^s-5-R¹⁹thiazole, and 2-Q^b-5-Q^s-4-R¹⁷thiazole;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, isopropyl, propyl, carboxy, amidino, guanidino, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl, ethylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, and cyano;

~~R¹⁸ or R¹⁹ is optionally C(NR²⁵)NR²³R²⁴ with the proviso that R¹⁸, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is C(NR²⁵)NR²³R²⁴ or hydrido, with the proviso that no more than one of R²³ and R²⁴ is hydroxy at the same time;

R²³, R²⁴, and R²⁵ are independently selected from the group consisting of hydrido, methyl, ethyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH₂ and CH₂CH₂.

11. (currently amended): The compound as recited in **[[Claim]] claim 10** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of 2-aminophenyl, 3-aminophenyl, 3-amidinophenyl, 4-amidinophenyl, 3-carboxyphenyl,

3-carboxy-5-hydroxyphenyl, 3-chlorophenyl, 4-chlorophenyl,
3,4-dichlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 3,4-difluorophenyl,
3-hydroxyphenyl, 4-hydroxyphenyl, 3-methoxyaminophenyl,
3-methoxyphenyl, 4-methoxyphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-
trifluoromethylphenyl, 2-imidazolyl, 2-pyridyl, 3-pyridyl,
5-chloro-3-trifluoromethyl-2-pyridyl, 4-pyridyl, 2-thienyl, 3-thienyl, and
3-trifluoromethyl-2-pyridyl;

A is selected from the group consisting of CH_2 , CH_3CH , CF_3CH , NHC(O) ,
 CH_2CH_2 , and $\text{CH}_2\text{CH}_2\text{CH}_2$;

M is N or $\text{R}^1\text{-C}$;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino,
hydroxyamino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy,
hydroxymethyl, methoxyamino, methylthio, trifluoromethoxy, fluoro, and chloro;



R^2 is $\text{Z}^0\text{-Q}$;

Z^0 is selected from the group consisting of a bond, O, S, NH, $\text{N}(\text{CH}_3)$, OCH_2 ,
and SCH_2 ;

Q is selected from the group consisting of
3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-
amino-5-benzylphenyl, 3-amino-5-(2-phenylethyl)phenyl,
3-amino-5-benzylaminophenyl, 3-amino-5-(2-phenylethylamino)phenyl,
3-amino-5-benzyloxyphenyl, 3-amino-5-(2-phenylethoxy)phenyl,
3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl,
3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl,
3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl,
3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl,
3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl,
3-amino-5-(N-benzylamidosulfonyl)phenyl,
3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl,
3-amino-5-(N-ethylamidocarbonyl)phenyl,
3-amino-5-(N-isopropylamidocarbonyl)phenyl,
3-amino-5-(N-propylamidocarbonyl)phenyl,
3-amino-5-(N-isobutylamidocarbonyl)phenyl,
3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl,
3-amino-5-(N-cyclobutylamidocarbonyl)phenyl,
3-amino-5-(N-cyclopentylamidocarbonyl)phenyl,

3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl, 3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl, 3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl, 3-aminophenyl, 3-amino-5-(4-trifluoromethylbenzylamino)phenyl, 3-amino-5-(4-trifluoromethylbenzyloxy)phenyl, 3-carboxyphenyl, 3-carboxy-5-hydroxyphenyl, 3-amino-5-carboxyphenyl, 3-chlorophenyl, 2-chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl, 2-fluorophenyl, 3-fluorophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 3-methanesulfonylaminophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 3-methoxyaminophenyl, 3-methoxycarbonylphenyl, 2-methylaminophenyl, 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl, 3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl, with the proviso that Q is other than a phenyl or substituted phenyl when Z⁰ is a bond;

Y⁰ is selected from the group consisting of:



~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹ benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹ pyridine, 3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹ pyridine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹ thiophene, and 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷ thiophene;~~

R^{16} and R^{19} are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano;

~~R^{16} or R^{19} is optionally $C(NR^{25})NR^{23}R^{24}$ with the proviso that R^{16} , R^{19} , and Q^b are not simultaneously hydrido;~~

R^{17} and R^{18} are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano;


Q^b is $C(NR^{25})NR^{23}R^{24}$ or hydrido;

R^{23} , R^{24} , and R^{25} are independently hydrido or methyl; **and**

Q^s is CH_2 .

Claims 12-16 (canceled).

17. (currently amended): The compound as recited in **[[Claim]] claim 2** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, C2-C8 alkyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} ;

R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b ;

A is a **bond** or $(\text{CH}(\text{R}^{15}))_{\text{pa}}-(\text{W}^7)_{\text{rr}}$ wherein rr is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is $(\text{R}^7)\text{NC}(\text{O})$ or $\text{N}(\text{R}^7)$;

R^7 is selected from the group consisting of hydrido, hydroxy and alkyl;

R^{15} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is N or $\text{R}^1\text{-C}$;

R^1 is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

R^2 is $\text{Z}^0\text{-Q}$;

Z^0 is a bond or $\text{W}^0\text{-(CH}(\text{R}^{42}))_{\text{p}}$ wherein p is 0 or 1 and W^0 is selected from the group consisting of O, S, and $\text{N}(\text{R}^{41})$;


R^{41} and R^{42} are independently hydrido or alkyl;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11}** ~~a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R^{13} , a carbon adjacent to R^9 and two atoms from the carbon at the point of attachment is optionally substituted by R^{10} , a carbon adjacent to R^{13} and two atoms from the carbon at the point of attachment is optionally substituted by R^{12} , and any carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than a phenyl when Z^0 is a bond;~~
 R^9 , R^{11} , and R^{13} are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkylsulfonamido,

alkylsulfinyl, alkylsulfonyl, amidosulfonyl, alkyl, alkoxy, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkoxyamino, alkylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, heterocyclylamino, heterocyclylalkylamino, alkylsulfonamido, amidosulfonyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, halo, haloalkyl, and cyano;

Y⁰ is **the formula**



wherein J⁵, J⁶, D⁵, D⁶ and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J⁵ and J⁶ is absent when T is a 5-membered heteroaryl ring, J⁵ is optionally substituted by R¹⁷ when J⁵ is a carbon atom, J⁶ is optionally substituted by R¹⁸ when J⁶ is a carbon atom, D⁵ is optionally substituted by R¹⁶ when D⁵ is a carbon atom and D⁶ is optionally substituted by R¹⁹ when D⁶ is a carbon atom;
~~phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s; a carbon two or three contiguous atoms from the point of attachment of Q^s to the phenyl or heteroaryl ring is substituted by Q^b; a carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁷; another carbon adjacent to the point of attachment of Q^s is optionally substituted by R¹⁸; a carbon adjacent to Q^b is optionally substituted by R¹⁶; and another carbon adjacent to Q^b is optionally substituted by R¹⁹;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino,

alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;

~~R¹⁶ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴),, with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that no more than one of R²⁰ and R²¹ is hydroxy ~~at the same time~~ and with the further proviso that no more than one of R²³ and R²⁴ is hydroxy ~~at the same time~~;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, alkyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH₂, and CH₂CH₂.

18. (currently amended): The compound as recited in **[[Claim]] claim 17** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propynyl, 2-propenyl, propyl, isopropyl, butyl, 2-butenyl, 3-butenyl, 2-butyryl, sec-butyl, tert-butyl, isobutyl, 2-methylpropenyl, 1-pentyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-pentyryl, 3-pentyryl, 2-pentyl, 1-methyl-2-butenyl, 1-methyl-3-butenyl, 1-methyl-2-butyryl, 3-pentyl, 1-ethyl-2-propenyl, 2-methylbutyl, 2-methyl-2-butenyl, 2-methyl-3-butenyl, 2-methyl-3-butyryl, 3-methylbutyl, 3-methyl-2-butenyl, 3-methyl-3-butenyl, 1-hexyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 2-hexyl, 1-methyl-2-pentenyl, 1-methyl-3-pentenyl, 1-methyl-4-pentenyl, 1-methyl-2-pentyryl, 1-methyl-3-pentyryl, 3-hexyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 1-propyl-2-propenyl, 1-ethyl-2-butyryl, 1-heptyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 6-heptenyl, 2-heptyryl, 3-heptyryl, 4-heptyryl, 5-heptyryl, 2-heptyl, 1-methyl-2-hexenyl, 1-methyl-3-hexenyl, 1-methyl-4-hexenyl, 1-methyl-5-hexenyl, 1-methyl-2-hexynyl, 1-methyl-3-hexynyl, 1-methyl-4-hexynyl, 3-heptyl, 1-ethyl-2-pentenyl, 1-ethyl-3-pentenyl, 1-ethyl-4-pentenyl, 1-butyl-2-propenyl, 1-ethyl-2-pentyryl, 1-ethyl-3-pentyryl, 2,2,2-trifluoroethyl, 2,2-difluoropropyl, 4-trifluoromethyl-5,5,5-trifluoropentyl, 4-trifluoromethylpentyl, 5,5,6,6,6-pentafluorohexyl, and 3,3,3-trifluoropropyl, wherein each member of group B is optionally substituted at any carbon up to and including 5 atoms from the point

of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} ;

R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, amidosulfonyl, N-methylamididosulfonyl, N,N-dimethylamididosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, cyano, and Q^b ;

A is selected from the group consisting of a bond, NH, $N(CH_3)$, $N(OH)$, CH_2 , CH_3CH , CF_3CH , $NHC(O)$, $N(CH_3)C(O)$, $C(O)NH$, $C(O)N(CH_3)$, CH_2CH_2 , $CH_2CH_2CH_2$, CH_3CHCH_2 , and CF_3CHCH_2 ;

M is N or R^1-C ;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, 1-aminoethyl, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, methoxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxyamino, methylthio, ethylthio, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, and bromo;

R^2 is Z^0-Q ;

Z^0 is selected from the group consisting of a bond, O, S, NH, $N(CH_3)$, OCH_2 , SCH_2 , $N(H)CH_2$, and $N(CH_3)CH_2$;

Q is selected from the group consisting of phenyl **[,]** and 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, and 1,3,5-triazin-2-yl **heteroaryl rings**, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative**

to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹ a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹; the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³; a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰; a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹²; and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond;




R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;





R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, carboxymethyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methanesulfonamido, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl,

N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidosulfonyl, N-(2-chlorobenzyl)amidosulfonyl, N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl, N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl, N-cyclohexylamidocarbonyl, fluoro, chloro, bromo, cyano, cyclobutoxy, cyclohexoxy, cyclohexylmethoxy, 4-trifluoromethylcyclohexylmethoxy, cyclopentoxyl, benzyl, benzyloxy, 4-bromo-3-fluorophenoxy, 3-bromobenzyloxy, 4-bromobenzyloxy, 4-bromobenzylamino, 5-bromopyrid-2-ylmethylamino, 4-butoxyphenamino, 3-chlorobenzyl, 4-chlorophenoxy, 4-chloro-3-ethylphenoxy, 4-chloro-3-ethylbenzylamino, 4-chloro-3-ethylphenylamino, 3-chlorobenzyloxy, 4-chlorobenzyloxy, 4-chlorobenzylsulfonyl, 4-chlorophenylamino, 4-chlorophenylsulfonyl, 5-chloropyrid-3-yloxy, 2-cyanopyrid-3-yloxy, 2,3-difluorobenzyloxy, 2,4-difluorobenzyloxy, 3,4-difluorobenzyloxy, 2,5-difluorobenzyloxy, 3,5-difluorophenoxy, 3,5-difluorobenzyloxy, 4-difluoromethoxybenzyloxy, 2,3-difluorophenoxy, 2,4-difluorophenoxy, 2,5-difluorophenoxy, 3,5-dimethylphenoxy, 3,4-dimethylphenoxy, 3,4-dimethylbenzyloxy, 3,5-dimethylbenzyloxy, 4-ethoxyphenoxy, 4-ethylbenzyloxy, 3-ethylphenoxy, 4-ethylaminophenoxy, 3-ethyl-5-methylphenoxy, 4-fluorobenzyloxy, 2-fluoro-3-trifluoromethylbenzyloxy, 3-fluoro-5-trifluoromethylbenzyloxy, 4-fluoro-2-trifluoromethylbenzyloxy, 4-fluoro-3-trifluoromethylbenzyloxy, 2-fluorophenoxy, 4-fluorophenoxy, 2-fluoro-3-trifluoromethylphenoxy, 2-fluorobenzyloxy, 4-fluorophenylamino, 2-fluoro-4-trifluoromethylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, phenylamino, 1-phenylethoxy, 2-phenylethoxy, 2-phenylethyl, 2-phenylethylamino, phenylsulfonyl, 3-trifluoromethoxybenzyloxy, 4-trifluoromethoxybenzyloxy, 3-trifluoromethoxyphenoxy, 4-trifluoromethoxyphenoxy, 3-trifluoromethylbenzyloxy, 4-trifluoromethylbenzyloxy, 2,4-bis-trifluoromethylbenzyloxy, 3-trifluoromethylbenzyl, 3,5-bis-trifluoromethylbenzyloxy, 4-trifluoromethylphenoxy, 3-trifluoromethylphenoxy, 3-trifluoromethylthiobenzyloxy,

4-trifluoromethylthiobenzyloxy, 2,3,4-trifluorophenoxy, 2,3,5-trifluorophenoxy, 3-pentafluoroethylphenoxy, 3-(1,1,2,2-tetrafluoroethoxy)phenoxy, and 3-trifluoromethylthiophenoxy;

Y⁰ is selected from the group consisting of:





~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹ benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹ pyridine,~~

~~3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridine, 2-Q^b-5-Q^s-3-R¹⁶-6-R¹⁸pyrazine, 3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹pyridazine, 2-Q^b-5-Q^s-4-R¹⁷-6-R¹⁸pyrimidine, 5-Q^b-2-Q^s-4-R¹⁶-6-R¹⁹pyrimidine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹thiophene, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷thiophene, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹furan, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷furan, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹pyrrole, 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷pyrrole, 4-Q^b-2-Q^s-5-R¹⁹imidazole, 2-Q^b-4-Q^s-5-R¹⁷imidazole, 3-Q^b-5-Q^s-4-R¹⁶isoxazole, 5-Q^b-3-Q^s-4-R¹⁶isoxazole, 2-Q^b-5-Q^s-4-R¹⁶pyrazole, 4-Q^b-2-Q^s-5-R¹⁹thiazole, and 2-Q^b-5-Q^s-4-R¹⁷thiazole;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, isopropyl, propyl, carboxy, amidino, guanidino, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl, ethylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, and cyano;

~~R¹⁶ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹; C(NR²⁵)NR²³R²⁴; and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, C(NR²⁵)NR²³R²⁴, and N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), with the proviso that no more than one of R²⁰ and R²¹ is hydroxy ~~at the same time~~ and with the further proviso that no more than one of R²³ and R²⁴ is hydroxy ~~at the same time~~;

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, methyl, ethyl, propyl, butyl, isopropyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH₂, and CH₂CH₂.

19. (currently amended): The compound as recited in **[[Claim]] claim 18** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butyl, (R)-2-butyl, (S)-2-butyl, tert-butyl, isobutyl, 1-pentyl, 3-pentyl, 2-methylbutyl, 2,2,2-trifluoroethyl, 6-amidocarbonylhexyl, 4-methyl-2-pentyl, 3-hydroxypropyl, 1-methoxy-2-propyl, 2-methoxyethyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 2-dimethylaminopropyl, 2-cyanoethyl, 6-hydroxyhexyl, 2-hydroxyethyl,

2-amidinoethyl, 2-guanidinoethyl, 3-guanidinopropyl, 4-guanidinobutyl, 3-hydroxypropyl, 4-hydroxybutyl, 6-cyanoethyl, 2-dimethylaminoethyl, 3-methylbutyl, 2-methylbutyl, (S)-2-methylbutyl, 3-aminopropyl, 2-hexyl, and 4-aminobutyl;

A is selected from the group consisting of a bond, CH_2 , NHC(O) , CH_2CH_2 , $\text{CH}_2\text{CH}_2\text{CH}_2$, and CH_3CHCH_2 ;

M is N or $\text{R}^1\text{-C}$;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy, hydroxymethyl, methoxyamino, methylthio, trifluoromethoxy, fluoro, and chloro;



R^2 is $\text{Z}^0\text{-Q}$;

Z^0 is selected from the group consisting of a bond, O, S, NH, $\text{N(CH}_3\text{)}$, OCH_2 , and SCH_2 ;

Q is selected from the group consisting of
3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-benzylphenyl, 3-amino-5-(2-phenylethyl)phenyl, 3-amino-5-benzylaminophenyl, 3-amino-5-(2-phenylethylamino)phenyl, 3-amino-5-benzyloxyphenyl, 3-amino-5-(2-phenylethoxy)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-benzylamidosulfonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, 3-amino-5-(N-ethylamidocarbonyl)phenyl, 3-amino-5-(N-isopropylamidocarbonyl)phenyl, 3-amino-5-(N-propylamidocarbonyl)phenyl, 3-amino-5-(N-isobutylamidocarbonyl)phenyl, 3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl, 3-amino-5-(N-cyclobutylamidocarbonyl)phenyl, 3-amino-5-(N-cyclopentylamidocarbonyl)phenyl, 3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl, 3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl, 3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl,

3-aminophenyl, 3-amino-5-(4-trifluoromethylbenzylamino)phenyl, 3-amino-5-(4-trifluoromethylbenzyloxy)phenyl, 3-carboxyphenyl, 3-carboxy-5-hydroxyphenyl, 3-amino-5-carboxyphenyl, 3-chlorophenyl, 2-chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl, 2-fluorophenyl, 3-fluorophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 3-methanesulfonylaminophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 3-methoxyaminophenyl, 3-methoxycarbonylphenyl, 2-methylaminophenyl, 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl, 3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl, with the proviso that Q is other than a phenyl or substituted phenyl when Z⁰ is a bond;

Y⁰ is selected from the group consisting of:



~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹ benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹ pyridine, 3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹ pyridine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹ thiophene, and 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷ thiophene;~~

R¹⁶ and R¹⁹ are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano;

~~R¹⁶ or R¹⁹ is optionally C(NR²⁵)NR²³R²⁴ with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;~~

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano;


Q^b is hydrido or C(NR²⁵)NR²³R²⁴;

R²³, R²⁴, and R²⁵ are independently hydrido or methyl; **and**

Q^s is CH₂.

Claims 20-24 (canceled).

25. (currently amended): The compound as recited in **[[Claim]] claim 2** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of C3-C7 cycloalkyl and C4 saturated heterocyclyl, wherein **(a) each ring carbon is optionally substituted with R³³, (b) a ring carbon, other than the ring carbon at the point of attachment, is optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, (c) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (d) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (e) a ring carbon or nitrogen, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (f) a ring carbon or nitrogen, if present, in a second beta**

position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², (g) a ring carbon or nitrogen, if present, in a first gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹⁰, is optionally substituted by R¹¹, and (h) a ring carbon or nitrogen, if present, in a second gamma position relative to the carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹², is optionally substituted by R³³; each ring carbon is optionally substituted with R³³; a ring carbon other than the ring carbon at the point of attachment of B to A is optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time; ring carbon sand a nitrogen adjacent to the carbon atom at the point of attachment is optionally substituted with R⁹ or R¹³; a ring carbon or nitrogen adjacent to the R⁹ position and two atoms from the point of attachment is optionally substituted with R¹⁰; a ring carbon or nitrogen adjacent to the R¹³ position and two atoms from the point of attachment is optionally substituted with R¹²; a ring carbon three atoms from the point of attachment and adjacent to the R¹⁰ position is optionally substituted with R¹¹; a ring carbon three atoms from the point of attachment and adjacent to the R¹² position is optionally substituted with R³³; and a ring carbon atoms from the point of attachment and adjacent to the R¹¹ and R³³ positions is optionally substituted with R³⁴;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkylsulfonamido, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, alkyl, alkoxy, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkoxyamino, alkylamino, arylamino, aralkylamino, heteroaryl amino, heteroaralkyl amino, heterocyclyl amino, heterocyclylalkyl amino, alkylsulfonamido, amidosulfonyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, halo, haloalkyl, and cyano;

R^{33} and R^{34} independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b ;

A is a bond or $(CH(R^{15}))_{pa}-(W^7)_r$ wherein rr is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is $(R^7)NC(O)$ or $N(R^7)$;

R^7 is selected from the group consisting of hydrido, hydroxy and alkyl;

R^{15} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is selected from the group consisting of N and R^1-C ;

R^1 is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

R^2 is Z^0-Q ;


Z^0 is a bond or $W^0-(CH(R^{42}))_p$ wherein p is 0 or 1 and W^0 is selected from the group consisting of O, S, and $N(R^{41})$;

R^{41} and R^{42} are independently hydrido or alkyl;

Q is phenyl or a heteroaryl of 5 or 6 ring members, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11}** ~~a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R^{13} , a carbon adjacent to R^9 and two atoms from the carbon at the point of attachment is optionally substituted by R^{10} , a carbon adjacent to R^{13} and two atoms from the carbon at the point of attachment is optionally~~

substituted by R^{12} , and any carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than a phenyl when Z^0 is a bond;

Y^0 is the formula



wherein J^5 , J^6 , D^5 , D^6 and the ring carbon atoms to which they are attached define a phenyl or 5- or 6-membered heteroaryl ring, T, wherein one of J^5 and J^6 is absent when T is a 5-membered heteroaryl ring, J^5 is optionally substituted by R^{17} when J^5 is a carbon atom, J^6 is optionally substituted by R^{18} when J^6 is a carbon atom, D^5 is optionally substituted by R^{16} when D^5 is a carbon atom and D^6 is optionally substituted by R^{19} when D^6 is a carbon atom;
~~phenyl or a heteroaryl of 5 or 6 ring members, wherein one carbon of said phenyl or said heteroaryl is substituted by Q^s , a carbon two or three contiguous atoms from the point of attachment of Q^s to the phenyl or heteroaryl ring is substituted by Q^b , a carbon adjacent to the point of attachment of Q^s is optionally substituted by R^{17} , another carbon adjacent to the point of attachment of Q^s is optionally substituted by R^{18} , a carbon adjacent to Q^b is optionally substituted by R^{16} , and another carbon adjacent to Q^b is optionally substituted by R^{19} ;~~

R^{16} , R^{17} , R^{18} , and R^{19} are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;

~~R^{16} or R^{19} is optionally $NR^{20}R^{21}$ or $C(NR^{25})NR^{23}R^{24}$, with the proviso that R^{16} , R^{19} , and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of $NR^{20}R^{21}$, hydrido, and $C(NR^{25})NR^{23}R^{24}$, with the proviso that no more than one of R^{20} and R^{21} is hydroxy at the same time and with the further proviso that no more than one of R^{23} and R^{24} is hydroxy at the same time;

R^{20} , R^{21} , R^{23} , R^{24} , and R^{25} are independently selected from the group consisting of hydrido, alkyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH_2 , and CH_2CH_2 .

26. (currently amended): The compound as recited in **[[Claim]] claim 25** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of cyclopropyl, cyclobutyl, oxetan-3-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, thiaetan-3-yl, cyclopentyl, cyclohexyl, norbornyl, 7-oxabicyclo[2.2.1]heptan-2-yl, bicyclo[3.1.0]hexan-6-yl, cycloheptyl, 2-morpholinyl, 3-morpholinyl, 4-morpholinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 2-dioxanyl, 4H-2-pyranyl, 4H-3-pyranyl, 4H-4-pyranyl, 4H-pyran-4-one-2-yl, 4H-pyran-4-one-3-yl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl, and 3-tetrahydrothienyl, wherein **(a) each ring carbon is optionally substituted with R³³, (b) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (c) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (d) a ring carbon or nitrogen, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, and (e) a ring carbon or nitrogen, if present, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹²;** ~~each ring carbon is optionally substituted with R³³; ring carbons and a nitrogen adjacent to the carbon atom at the point of attachment are optionally substituted with R⁹ or R¹³; a ring carbon or nitrogen adjacent to the R⁹ position and two atoms from the point of attachment is optionally substituted with R¹⁰; and a ring carbon or nitrogen adjacent to the R¹³ position and two atoms from the point of attachment is optionally substituted with R¹²;~~

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, methanesulfonamido, amidosulfonyl, N-methylamidulosulfonyl, N,N-

dimethylamidodisulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, carboxymethyl, methyl, ethyl, propyl, isopropyl, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methanesulfonamido, amidodisulfonyl, N-methylamidodisulfonyl, N,N-dimethylamidodisulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl, N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidodisulfonyl, N-(2-chlorobenzyl)amidodisulfonyl, N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl, N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl, N-cyclohexylamidocarbonyl, fluoro, chloro, bromo, cyano, cyclobutoxy, cyclohexoxy, cyclohexylmethoxy, 4-trifluoromethylcyclohexylmethoxy, cyclopentoxy, benzyl, benzyloxy, 4-bromo-3-fluorophenoxy, 3-bromobenzyloxy, 4-bromobenzyloxy, 4-bromobenzylamino, 5-bromopyrid-2-ylmethylamino, 4-butoxyphenamino, 3-chlorobenzyl, 4-chlorophenoxy, 4-chloro-3-ethylphenoxy, 4-chloro-3-ethylbenzylamino, 4-chloro-3-ethylphenylamino, 3-chlorobenzyloxy, 4-chlorobenzyloxy, 4-chlorobenzylsulfonyl, 4-chlorophenylamino, 4-chlorophenylsulfonyl, 5-chloropyrid-3-yl, 2-cyanopyrid-3-yl, 2,3-difluorobenzyloxy, 2,4-difluorobenzyloxy, 3,4-difluorobenzyloxy, 2,5-difluorobenzyloxy, 3,5-difluorophenoxy, 3,5-difluorobenzyloxy, 4-difluoromethoxybenzyloxy, 2,3-difluorophenoxy, 2,4-difluorophenoxy, 2,5-difluorophenoxy, 3,5-dimethylphenoxy, 3,4-dimethylphenoxy, 3,4-dimethylbenzyloxy, 3,5-dimethylbenzyloxy, 4-ethoxyphenoxy, 4-ethylbenzyloxy, 3-ethylphenoxy, 4-ethylaminophenoxy, 3-ethyl-5-methylphenoxy, 4-fluorobenzyloxy, 2-fluoro-3-trifluoromethylbenzyloxy, 3-fluoro-5-trifluoromethylbenzyloxy,

4-fluoro-2-trifluoromethylbenzyloxy, 4-fluoro-3-trifluoromethylbenzyloxy, 2-fluorophenoxy, 4-fluorophenoxy, 2-fluoro-3-trifluoromethylphenoxy, 2-fluorobenzyloxy, 4-fluorophenylamino, 2-fluoro-4-trifluoromethylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, 4-isopropylbenzyloxy, 3-isopropylphenoxy, 4-isopropylphenoxy, 4-isopropyl-3-methylphenoxy, phenylamino, 1-phenylethoxy, 2-phenylethoxy, 2-phenylethyl, 2-phenylethylamino, phenylsulfonyl, 3-trifluoromethoxybenzyloxy, 4-trifluoromethoxybenzyloxy, 3-trifluoromethoxyphenoxy, 4-trifluoromethoxyphenoxy, 3-trifluoromethylbenzyloxy, 4-trifluoromethylbenzyloxy, 2,4-bis-trifluoromethylbenzyloxy, 3-trifluoromethylbenzyl, 3,5-bis-trifluoromethylbenzyloxy, 4-trifluoromethylphenoxy, 3-trifluoromethylphenoxy, 3-trifluoromethylthiobenzyloxy, 4-trifluoromethylthiobenzyloxy, 2,3,4-trifluorophenoxy, 2,3,5-trifluorophenoxy, 3-pentafluoroethylphenoxy, 3-(1,1,2,2-tetrafluoroethoxy)phenoxy, and 3-trifluoromethylthiophenoxy;

R^{33} and R^{34} are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, methoxy, ethoxy, isopropoxy, propoxy, hydroxy, amino, methoxyamino, ethoxyamino, acetamido, trifluoroacetamido, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, amidosulfonyl, N-methylamidodisulfonyl, N,N-dimethylamidodisulfonyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2,2,2-trifluoro-1-hydroxyethyl, methoxycarbonyl, ethoxycarbonyl, amidocarbonyl, N-methylamidocarbonyl, N,N-dimethylamidocarbonyl, cyano, and Q^b ;

A is selected from the group consisting of a bond, NH, $N(CH_3)$, $N(OH)$, CH_2 , CH_3CH , CF_3CH , $NHC(O)$, $N(CH_3)C(O)$, $C(O)NH$, $C(O)N(CH_3)$, CH_2CH_2 , $CH_2CH_2CH_2$, CH_3CHCH_2 , and CF_3CHCH_2 ;

M is N or R^1-C ;

R^1 is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, 1-aminoethyl, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, methoxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, methoxyamino, methylthio, ethylthio, trifluoromethoxy,





1,1,2,2-tetrafluoroethoxy, fluoro, chloro, and bromo;




R^2 is Z^0 -Q;

Z^0 is selected from the group consisting of a bond, O, S, NH, $N(CH_3)$, OCH_2 , SCH_2 , $N(H)CH_2$, and $N(CH_3)CH_2$;

Q is selected from the group consisting of phenyl **[[,]]** **and** 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, 5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, and 1,3,5-triazin-2-yl **heteroaryl rings**, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11}** a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z^0 is optionally substituted by R^9 , the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R^{13} , a carbon adjacent to R^9 and two atoms from the carbon at the point of attachment is optionally substituted by R^{10} , a carbon adjacent to R^{13} and two atoms from the carbon at the point of attachment is optionally substituted by R^{12} , and any carbon adjacent to both R^{10} and R^{12} is optionally substituted by R^{11} , with the proviso that Q is other than a phenyl when Z^0 is a bond;

Y^0 is selected from the group consisting of:





~~1-Q^b-4-Q^s-2-R¹⁸-3-R¹⁷-5-R¹⁸-6-R¹⁹benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹pyridine, 3-Q^b-6-Q^s-2-R¹⁸-5-R¹⁸-4-R¹⁹pyridine, 2-Q^b-5-Q^s-3-R¹⁸-6-R¹⁸pyrazine, 3-Q^b-6-Q^s-2-R¹⁸-5-R¹⁸-4-R¹⁹pyridazine, 2-Q^b-5-Q^s-4-R¹⁷-6-R¹⁸pyrimidine, 5-Q^b-2-Q^s-4-R¹⁸-6-R¹⁹pyrimidine, 3-Q^b-5-Q^s-4-R¹⁸-2-R¹⁹thiophene, 2-Q^b-5-Q^s-3-R¹⁸-4-R¹⁷thiophene, 3-Q^b-5-Q^s-4-R¹⁸-2-R¹⁹furan, 2-Q^b-5-Q^s-3-R¹⁸-4-R¹⁷furan, 3-Q^b-5-Q^s-4-R¹⁸-2-R¹⁹pyrrole, 2-Q^b-5-Q^s-3-R¹⁸-4-R¹⁷pyrrole, 4-Q^b-2-Q^s-5-R¹⁹imidazole, 2-Q^b-4-Q^s-5-R¹⁷imidazole, 3-Q^b-5-Q^s-4-R¹⁸isoxazole, 5-Q^b-3-Q^s-4-R¹⁸isoxazole, 2-Q^b-5-Q^s-4-R¹⁸pyrazole, 4-Q^b-2-Q^s-5-R¹⁹thiazole, and 2-Q^b-5-Q^s-4-R¹⁷thiazole;~~

R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, methyl, ethyl, isopropyl, propyl, carboxy, amidino, guanidino, methoxy,

ethoxy, isopropoxy, propoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, N-ethylamino, methylthio, ethylthio, isopropylthio, trifluoromethylthio, methylsulfinyl, ethylsulfinyl, methylsulfonyl, ethylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, 2,2,3,3,3-pentafluoropropyl, trifluoromethoxy, 1,1,2,2-tetrafluoroethoxy, fluoro, chloro, bromo, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, and cyano;

~~R¹⁶ or R¹⁹ is optionally C(NR²⁵)NR²³R²⁴ with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is C(NR²⁵)NR²³R²⁴ or hydrido, with the proviso that no more than one of R²³ and R²⁴ is hydroxy ~~at the same time;~~

R²³, R²⁴, and R²⁵ are independently selected from the group consisting of hydrido, methyl, ethyl, and hydroxy; **and**

Q^s is selected from the group consisting of a bond, CH₂ and CH₂CH₂.

27. (currently amended): The compound as recited in **[[Claim]] claim 26** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, oxalan-2-yl, 2-(2R)-bicyclo[2.2.1]-heptyl, 1-pyrrolidinyl, 1-piperidinyl, oxetan-3-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, 7-oxabicyclo[2.2.1]heptan-2-yl, bicyclo[3.1.0]hexan-6-yl, 2-morpholinyl, 3-morpholinyl, 4-morpholinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 2-dioxanyl, 4H-2-pyranyl, 4H-3-pyranyl, 4H-4-pyranyl, 4H-pyran-4-one-2-yl, 4H-pyran-4-one-3-yl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl, and 3-tetrahydrothienyl;

A is selected from the group consisting of a bond, CH₂, NHC(O), CH₂CH₂, and CH₂CH₂CH₂;

M is selected from the group consisting of N and R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, amino, amidino, hydroxyamino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy, hydroxymethyl, methoxyamino, methylthio, trifluoromethoxy, fluoro, and chloro;



R² is Z⁰-Q;

Z⁰ is selected from the group consisting of a bond, O, S, NH, N(CH₃), OCH₂, and SCH₂;

Q is selected from the group consisting of
3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-benzylphenyl, 3-amino-5-(2-phenylethyl)phenyl, 3-amino-5-benzylaminophenyl, 3-amino-5-(2-phenylethylamino)phenyl, 3-amino-5-benzyloxyphenyl, 3-amino-5-(2-phenylethoxy)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-benzylamidosulfonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, 3-amino-5-(N-ethylamidocarbonyl)phenyl, 3-amino-5-(N-isopropylamidocarbonyl)phenyl, 3-amino-5-(N-propylamidocarbonyl)phenyl, 3-amino-5-(N-isobutylamidocarbonyl)phenyl, 3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl, 3-amino-5-(N-cyclobutylamidocarbonyl)phenyl, 3-amino-5-(N-cyclopentylamidocarbonyl)phenyl, 3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl, 3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl, 3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl, 3-aminophenyl, 3-amino-5-(4-trifluoromethylbenzylamino)phenyl, 3-amino-5-(4-trifluoromethylbenzyloxy)phenyl, 3-carboxyphenyl, 3-carboxy-5-hydroxyphenyl, 3-amino-5-carboxyphenyl, 3-chlorophenyl, 2-chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl, 2-fluorophenyl, 3-fluorophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 3-methanesulfonylaminophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 3-methoxyaminophenyl, 3-methoxycarbonylphenyl, 2-methylaminophenyl, 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl,

3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl, with the proviso that Q is other than a phenyl or substituted phenyl when Z⁰ is a bond;

Y⁰ is selected from the group consisting of:



~~1-Q^b-4-Q^s-2-R¹⁶-3-R¹⁷-5-R¹⁸-6-R¹⁹ benzene, 2-Q^b-5-Q^s-6-R¹⁷-4-R¹⁸-3-R¹⁹ pyridine, 3-Q^b-6-Q^s-2-R¹⁶-5-R¹⁸-4-R¹⁹ pyridine, 3-Q^b-5-Q^s-4-R¹⁶-2-R¹⁹ thiophene, and 2-Q^b-5-Q^s-3-R¹⁶-4-R¹⁷ thiophene;~~

R¹⁶ and R¹⁹ are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano;

~~R¹⁶ or R¹⁹ is optionally C(NR²⁵)NR²³R²⁴ with the proviso that R¹⁶, R¹⁹, and Q^b are not simultaneously hydrido;~~

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano;


Q^b is C(NR²⁵)NR²³R²⁴ or hydrido;

R²³, R²⁴, and R²⁵ are independently hydrido or methyl; **and**

Q^s is CH₂.

Claims 28-32 (canceled).

33. (currently amended): Compound of ~~[[Claim]]~~ **claim** 2 of the Formula:



or a

pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of:

(i) ~~[[B is]] phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³², (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³⁶, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³², is optionally substituted by R³³, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³⁶, is optionally substituted by R³⁵, and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R³³ and R³⁵, respectively, is optionally substituted by R³⁴; a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³²; the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R³⁶; a carbon adjacent to R³² and two atoms from the carbon at the point of attachment is optionally substituted by R³³; a carbon adjacent to R³⁶ and two atoms from the carbon at the point of attachment is optionally substituted by R³⁵; and any carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;~~

(ii) hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to 6 atoms from the point of

attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶; and

(iii) a C3-C12 cycloalkyl or a C4-C9 saturated heterocyclyl, wherein (a) each ring carbon may be optionally substituted with R₃₃, (b) a ring carbon, other than the ring carbon at the point of attachment of B to A, may be optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, (c) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (d) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (e) a ring carbon or nitrogen atom, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (f) a ring carbon or nitrogen, if present, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², (g) a ring carbon or nitrogen, if present, in a first gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹⁰, is optionally substituted by R¹¹, (h) a ring carbon or nitrogen, if present, in a second gamma position relative to the carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹², is optionally substituted by R³³, and (i) a ring carbon or nitrogen, if present, in a delta position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹¹ and R³³, respectively, is optionally substituted by R³⁴;

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkylenedioxy, haloalkylthio, alkanoyloxy, alkoxy, hydroxy, amino, alkoxyamino, haloalkanoyl, nitro, alkylamino, alkylthio, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, alkylsulfonamido, amidosulfonyl, alkyl, alkenyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyalkyl, alkylamino, carboalkoxy, carboxy, carboxamido, cyano, and Q^b;

~~— B is optionally selected from the group consisting of hydrido, trialkylsilyl, C2-C8 alkyl, C3-C8 alkylenyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl;~~

wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} ;

~~B is optionally a C3-C12 cycloalkyl or a C4-C9 saturated heterocyclyl, wherein each ring carbon is optionally substituted with R^{33} , a ring carbon other than the ring carbon at the point of attachment of B to A is optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, ring carbons and a nitrogen adjacent to the carbon atom at the point of attachment are optionally substituted with R^9 or R^{13} , a ring carbon or nitrogen adjacent to the R^9 position and two atoms from the point of attachment is optionally substituted with R^{10} , a ring carbon or nitrogen adjacent to the R^{13} position and two atoms from the point of attachment is optionally substituted with R^{12} , a ring carbon or nitrogen three atoms from the point of attachment and adjacent to the R^{10} position is optionally substituted with R^{11} , a ring carbon or nitrogen three atoms from the point of attachment and adjacent to the R^{12} position is optionally substituted with R^{33} , and a ring carbon or nitrogen four atoms from the point of attachment and adjacent to the R^{11} and R^{33} positions is optionally substituted with R^{34} ;~~

R^9 , R^{10} , R^{11} , R^{12} , and R^{13} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, alkoxyamino, alkanoyl, haloalkanoyl, amidino, guanidino, alkylenedioxy, haloalkylthio, alkoxy, cycloalkoxy, cycloalkylalkoxy, aralkoxy, aryloxy, heteroaryloxy, heteroaralkoxy, heterocyclyloxy, heterocyclylalkoxy, hydroxy, amino, alkylamino, N-alkyl-N-aryl amino, arylamino, aralkylamino, heteroaryl amino, heteroaralkyl amino, heterocyclyl amino, heterocyclylalkyl amino, alkylthio, alkylsulfinyl, arylsulfinyl, aralkylsulfinyl, cycloalkylsulfinyl, heteroarylsulfinyl, alkylsulfamido, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl, cycloalkylsulfonyl, heteroarylsulfonyl, amidosulfonyl, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heterocyclyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, hydroxyhaloalkyl, aminoalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxamido, and cyano;

A is a bond or $(CH(R^{15}))_{pa}-(W^7)_r$ wherein r is 0 or 1, pa is an integer selected from 0 through 3, and W^7 is selected from the group consisting of O, S, C(O), $(R^7)NC(O)$, $(R^7)NC(S)$, and $N(R^7)$;

R^7 is selected from the group consisting of hydrido, hydroxy and alkyl;

R^{15} is selected from the group consisting of hydrido, hydroxy, halo, alkyl, and haloalkyl;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, alkyl, cyano, halo, haloalkyl, haloalkoxy, amino, aminoalkyl, alkylamino, amidino, hydroxy, hydroxyamino, alkoxy, hydroxyalkyl, alkoxyamino, thiol, and alkylthio;

R² is Z⁰-Q;

Z⁰ is selected from the group consisting of:

(i) a bond, W⁰-(CH(R⁴²))_p wherein p is an integer selected from 0 through 3 and W⁰ is selected from the group consisting of O, S, and N(R⁴¹), and (CH(R⁴¹))_g-O wherein g is an integer selected from 1 through 3, with the proviso that Z⁰ is directly bonded to the pyrimidinone ring; **and**

(ii) ~~Z⁰ is optionally~~ W²²-(CH(R⁴²))_h wherein h is 0 or 1 and W²² is selected from the group consisting of 1,2-cyclopropyl, 1,2-cyclobutyl, 1,2-cyclohexyl, 1,3-cyclohexyl, 1,2-cyclopentyl, 1,3-cyclopentyl, 2,3-morpholinyl, 2,4-morpholinyl, 2,6-morpholinyl, 3,4-morpholinyl, 3,5-morpholinyl, 1,2-piperazinyl, 1,3-piperazinyl, 2,3-piperazinyl, 2,6-piperazinyl, 1,2-piperidinyl, 1,3-piperidinyl, 2,3-piperidinyl, 2,4-piperidinyl, 2,6-piperidinyl, 3,4-piperidinyl, 1,2-pyrrolidinyl, 1,3-pyrrolidinyl, 2,3-pyrrolidinyl, 2,4-pyrrolidinyl, 2,5-pyrrolidinyl, 3,4-pyrrolidinyl, 2,3-tetrahydrofuranyl, 2,4-tetrahydrofuranyl, 2,5-tetrahydrofuranyl, and 3,4-tetrahydrofuranyl, wherein Z⁰ is directly bonded to the pyrimidinone ring and W²² is optionally substituted with one or more substituents selected from the group consisting of R⁹, R¹⁰, R¹¹, R¹², and R¹³;

R⁴¹ and R⁴² are independently selected from the group consisting of hydrido, hydroxy, and amino;

Q is selected from the group consisting of:

(i) **[[Q is]] phenyl or a heteroaryl of 5 or 6 ring members, wherein (a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, if**

present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹ a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond; **and**

(ii) Q is optionally hydrido with the proviso that Z⁰ is selected from other than a bond;

K is CHR^{4a} wherein R^{4a} is selected from the group consisting of hydrido, hydroxyalkyl, alkyl, alkoxyalkyl, alkylthioalkyl, and haloalkyl;

E⁰ is selected from the group consisting of a bond, C(O)N(H), (H)NC(O), (R⁷)NS(O)₂, and S(O)₂N(R⁷);

Y^{AT} is Q^b-Q^s;

Q^s is (CR³⁷R³⁸)_b wherein b is an integer selected from 1 through 4, R³⁷ is selected from the group consisting of hydrido, alkyl, and haloalkyl, and R³⁸ is selected from the group consisting of hydrido, alkyl, haloalkyl, aroyl, and heteroaroyl with the proviso that there is at least one aroyl or heteroaroyl substituent, with the further proviso that no more than one aroyl or heteroaroyl is bonded to (CR³⁷R³⁸)_b at the same time, with the still further proviso that said aroyl and said heteroaroyl are optionally substituted with one or more substituents selected from the group consisting of R¹⁶, R¹⁷, R¹⁸, and R¹⁹, with another further proviso that said aroyl and said heteroaroyl are bonded to the CR³⁷R³⁸ that is directly bonded to E⁰, with still another further proviso that no more than one alkyl or one haloalkyl is bonded to a CR³⁷R³⁸ at the same time, and with the additional proviso that said alkyl and haloalkyl are bonded to a carbon other than the one bonding said aroyl or said heteroaroyl;


R¹⁶, R¹⁷, R¹⁸, and R¹⁹ are independently selected from the group consisting of hydrido, amidino, guanidino, carboxy, haloalkylthio, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, alkanoyl, haloalkanoyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, aminoalkyl, and cyano;

~~R¹⁰ or R¹⁹ is optionally selected from the group consisting of NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that R¹⁰, R¹⁹, and Q^b are not simultaneously hydrido;~~

Q^b is selected from the group consisting of NR²⁰R²¹, hydrido, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴, with the proviso that no more than one of R²⁰ and R²¹ is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino ~~at the same time~~ and with the further proviso that no more than one of R²³ and R²⁴ is selected from the group consisting of hydroxy, amino, alkylamino, and dialkylamino ~~at the same time~~; **and**

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido, alkyl, hydroxy, amino, alkylamino and dialkylamino.

34. (currently amended): The compound as recited in **[[Claim]] claim 33** having the Formula:



or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of:

~~(i) B is selected from the group consisting of phenyl **[[,]]** and 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 3-isoxazolyl, and 5-isoxazolyl heteroaryl rings, wherein~~
(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³², (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R³⁶, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R³², is optionally substituted by R³³, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom

optionally substituted by R³⁶, is optionally substituted by R³⁵, and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R³³ and R³⁵, respectively, is optionally substituted by R³⁴; a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to A is optionally substituted by R³²; the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R³⁶; a carbon adjacent to R³² and two atoms from the carbon at the point of attachment is optionally substituted by R³³; a carbon adjacent to R³⁶ and two atoms from the carbon at the point of attachment is optionally substituted by R³⁵; and any carbon adjacent to both R³³ and R³⁵ is optionally substituted by R³⁴;

(ii) hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butenyl, 2-butylnyl, sec-butyl, tert-butyl, isobutyl, 2-methylpropenyl, 1-pentyl, 2-pentenyl, 3-pentenyl, 2-pentylnyl, 3-pentylnyl, 2-pentyl, 3-pentyl, 2-methylbutyl, 2-methyl-2-butenyl, 3-methylbutyl, 3-methyl-2-butenyl, 1-hexyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 2-hexyl, 1-methyl-2-pentenyl, 1-methyl-3-pentenyl, 1-methyl-2-pentylnyl, 1-methyl-3-pentylnyl, 3-hexyl, 1-ethyl-2-butenyl, 1-heptyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 2-heptylnyl, 3-heptylnyl, 4-heptylnyl, 5-heptylnyl, 2-heptyl, 1-methyl-2-hexenyl, 1-methyl-3-hexenyl, 1-methyl-4-hexenyl, 1-methyl-2-hexynyl, 1-methyl-3-hexynyl, 1-methyl-4-hexynyl, 3-heptyl, 1-ethyl-2-pentenyl, 1-ethyl-3-pentenyl, 1-ethyl-2-pentylnyl, 1-ethyl-3-pentylnyl, 2,2,2-trifluoroethyl, 2,2-difluoropropyl, 4-trifluoromethyl-5,5,5-trifluoropentyl, 4-trifluoromethylpentyl, 5,5,6,6,6-pentafluorohexyl, and 3,3,3-trifluoropropyl, wherein each member of group B is optionally substituted at any carbon up to 5 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶; and

(iii) cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, oxalan-2-yl, 2-(2R)-bicyclo[2.2.1]-heptyl, oxetan-3-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, bicyclo[3.1.0]hexan-6-yl, 2-morpholinyl, 3-morpholinyl, 4-morpholinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 2-dioxanyl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl, and 3-tetrahydrothienyl, wherein (a) each ring carbon is optionally substituted with

R³³, (b) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (c) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (d) a ring carbon or nitrogen, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, and (e) a ring carbon or nitrogen, if present, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹²:

R³², R³³, R³⁴, R³⁵, and R³⁶ are independently selected from the group consisting of hydrido, amidino, guanidino, methyl, ethyl, methoxy, ethoxy, hydroxy, amino, N-methylamino, dimethylamino, methoxyamino, methylthio, ethylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, fluoro, chloro, bromo, amidosulfonyl, N-methylamidosulfonyl, hydroxymethyl, amidocarbonyl, carboxy, cyano, and Q^b;

~~B is optionally selected from the group consisting of hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butenyl, 2-butylnyl, sec-butyl, tert-butyl, isobutyl, 2-methylpropenyl, 1-pentyl, 2-pentenyl, 3-pentenyl, 2-pentylnyl, 3-pentylnyl, 2-pentyl, 3-pentyl, 2-methylbutyl, 2-methyl-2-butenyl, 3-methylbutyl, 3-methyl-2-butenyl, 1-hexyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 2-hexyl, 1-methyl-2-pentenyl, 1-methyl-3-pentenyl, 1-methyl-2-pentylnyl, 1-methyl-3-pentylnyl, 3-hexyl, 1-ethyl-2-butenyl, 1-heptyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 5-heptenyl, 2-heptylnyl, 3-heptylnyl, 4-heptylnyl, 5-heptylnyl, 2-heptyl, 1-methyl-2-hexenyl, 1-methyl-3-hexenyl, 1-methyl-4-hexenyl, 1-methyl-2-hexynyl, 1-methyl-3-hexynyl, 1-methyl-4-hexynyl, 3-heptyl, 1-ethyl-2-pentenyl, 1-ethyl-3-pentenyl, 1-ethyl-2-pentylnyl, 1-ethyl-3-pentylnyl, 2,2,2-trifluoroethyl, 2,2-difluoropropyl, 4-trifluoromethyl-5,5,5-trifluoropentyl, 4-trifluoromethylpentyl, 5,5,6,6,6-pentafluorohexyl, and 3,3,3-trifluoropropyl, wherein each member of group B is optionally substituted at any carbon up to and including 5 atoms from the point of attachment of B to A with one or more of the group consisting of R³², R³³, R³⁴, R³⁵, and R³⁶;~~

~~— B is optionally selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, oxalan-2-yl, 2-(2R)-bicyclo[2.2.1]-heptyl, oxetan-3-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, bicyclo[3.1.0]hexan-6-yl, 2-morpholinyl, 3-morpholinyl, 4-morpholinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 2-dioxanyl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl, and 3-tetrahydrothienyl, wherein each ring carbon is optionally substituted with R³³, ring carbons and a nitrogen adjacent to the carbon atom at the point of attachment are optionally substituted with R⁹ or R¹³, a ring carbon or nitrogen adjacent to the R⁹ position and two atoms from the point of attachment is optionally substituted with R¹⁰, and a ring carbon or nitrogen adjacent to the R¹³ position and two atoms from the point of attachment is optionally substituted with R¹²;~~

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, methyl, ethyl, methoxy, ethoxy, hydroxy, amino, N-methylamino, N,N-dimethylamino, methylthio, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, fluoro, chloro, bromo, amidosulfonyl, N-methylamidosulfonyl, N,N-dimethylamidosulfonyl, hydroxymethyl, 1-hydroxyethyl, amidocarbonyl, N-methylamidocarbonyl, carboxy, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, amidino, amidocarbonyl, N-methylamidocarbonyl, N-benzylamidocarbonyl, N-(2-chlorobenzyl)amidocarbonyl, N-(3-fluorobenzyl)amidocarbonyl, N-(2-trifluoromethylbenzyl)amidocarbonyl, N-(1-phenylethyl)amidocarbonyl, N-(1-methyl-1-phenylethyl)amidocarbonyl, N-benzylamidosulfonyl, N-(2-chlorobenzyl)amidosulfonyl, N-ethylamidocarbonyl, N-isopropylamidocarbonyl, N-propylamidocarbonyl, N-isobutylamidocarbonyl, N-(2-butyl)amidocarbonyl, N-cyclobutylamidocarbonyl, N-cyclopentylamidocarbonyl, N-cyclohexylamidocarbonyl, guanidino, methyl, ethyl, methoxy, ethoxy, hydroxy, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, carboxy, carboxymethyl, amino, acetamido, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, trifluoroacetamido, aminomethyl, N-methylamino, dimethylamino, methoxyamino, amidosulfonyl, N-methylamidosulfonyl,

N,N-dimethylamidodisulfonyl, methanesulfonamido, methoxycarbonyl, fluoro, chloro, bromo, and cyano;

A is selected from the group consisting of a bond, NH, N(CH₃), CH₂, CH₃CH, CH₂CH₂, and CH₂CH₂CH₂;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxymethyl, amino, aminomethyl, methylamino, cyano, methyl, trifluoromethyl, methoxy, methylthio, trifluoromethoxy, fluoro, and chloro;

R² is Z⁰-Q;

Z⁰ is selected from the group consisting of a bond, O, S, NH, OCH₂, SCH₂, and N(H)CH₂;

Q is selected from the group consisting of phenyl **[,]** **and** 2-thienyl, 2-furyl, 2-pyrrolyl, 2-imidazolyl, 2-thiazolyl, 3-isoxazolyl, 2-pyridyl, and 3-pyridyl **heteroaryl rings**, wherein **(a) a ring carbon in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹** ~~a carbon adjacent to the carbon at the point of attachment of said phenyl or heteroaryl ring to Z⁰ is optionally substituted by R⁹, the other carbon adjacent to the carbon at the point of attachment is optionally substituted by R¹³, a carbon adjacent to R⁹ and two atoms from the carbon at the point of attachment is optionally substituted by R¹⁰, a carbon adjacent to R¹³ and two atoms from the carbon at the point of attachment is optionally substituted by R¹², and any carbon adjacent to both R¹⁰ and R¹² is optionally substituted by R¹¹, with the proviso that Q is other than a phenyl when Z⁰ is a bond;~~

Y^{AT} is Q^b-Q^s;

Q^s is selected from the group consisting of: C[R³⁷(benzoyl)(CR³⁷R³⁸)_b],

$C[R^{37}(2\text{-pyridylcarbonyl})(CR^{37}R^{38})_b]$,
 $C[R^{37}(3\text{-pyridylcarbonyl})(CR^{37}R^{38})_b]$,
 $C[R^{37}(4\text{-pyridylcarbonyl})(CR^{37}R^{38})_b]$,
 $C[R^{37}(2\text{-thienylcarbonyl})(CR^{37}R^{38})_b]$,
 $C[R^{37}(3\text{-thienylcarbonyl})(CR^{37}R^{38})_b]$,
 $C[R^{37}(2\text{-thiazolylcarbonyl})(CR^{37}R^{38})_b]$,
 $C[R^{37}(4\text{-thiazolylcarbonyl})(CR^{37}R^{38})_b]$, and
 $C[R^{37}(5\text{-thiazolylcarbonyl})(CR^{37}R^{38})_b]$, wherein b is an integer selected from 1 through 3, R^{37} and R^{38} are independently selected from the group consisting of hydrido, alkyl, and haloalkyl, with the proviso that said benzoyl and the heteroaroyls are optionally substituted with one or more substituents selected from the group consisting of R^{16} , R^{17} , R^{18} , and R^{19} with the proviso that R^{17} and R^{18} are optionally substituted at a carbon selected from other than the meta and para carbons relative to the carbonyl of the benzoyl or heteroaroyl, with the further proviso that said benzoyl or said heteroaroyl are bonded to the carbon directly bonded to amide nitrogen of the 1-(amidocarbonylmethylene) group, and with the still further proviso that is no more than one alkyl or one haloalkyl is bonded to a $CR^{37}R^{38}$ at the same time;

R^{16} , R^{17} , R^{18} , and R^{19} are independently selected from the group consisting of hydrido, methyl, ethyl, amidino, guanidino, methoxy, hydroxy, amino, aminomethyl, 1-aminoethyl, 2-aminoethyl, N-methylamino, dimethylamino, methylthio, ethylthio, trifluoromethylthio, methylsulfinyl, methylsulfonyl, trifluoromethyl, pentafluoroethyl, 2,2,2-trifluoroethyl, trifluoromethoxy, fluoro, chloro, hydroxymethyl, carboxy, and cyano;

Q^b is $C(NR^{25})NR^{23}R^{24}$ or $N(R^{26})C(NR^{25})N(R^{23})(R^{24})$; **and**

R^{23} , R^{24} , R^{25} , and R^{26} are independently selected from the group consisting of hydrido, methyl, and ethyl.

35. (currently amended): The compound as recited in **[[Claim]] claim 34** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of:

(i) 2-aminophenyl, 3-aminophenyl, 3-amidinophenyl, 4-amidinophenyl, 3-carboxyphenyl, 3-carboxy-5-hydroxyphenyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-dichlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 3,4-difluorophenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 3-methoxyaminophenyl,

3-methoxyphenyl, 4-methoxyphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-trifluoromethylphenyl, 2-imidazolyl, 2-pyridyl, 3-pyridyl, 5-chloro-3-trifluoromethyl-2-pyridyl, 4-pyridyl, 2-thienyl, 3-thienyl, and 3-trifluoromethyl-2-pyridyl;

~~(ii) B is optionally selected from the group consisting of~~ hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butyl, (R)-2-butyl, (S)-2-butyl, tert-butyl, isobutyl, 1-pentyl, 3-pentyl, 2-methylbutyl, 2,2,2-trifluoroethyl, 6-amidocarbonylhexyl, 4-methyl-2-pentyl, 3-hydroxypropyl, 1-methoxy-2-propyl, 2-methoxyethyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 2-dimethylaminopropyl, 2-cyanoethyl, 6-hydroxyhexyl, 2-hydroxyethyl, 2-amidinoethyl, 2-guanidinoethyl, 3-guanidinopropyl, 4-guanidinobutyl, 3-hydroxypropyl, 4-hydroxybutyl, 6-cyanoethyl, 2-dimethylaminoethyl, 3-methylbutyl, 2-methylbutyl, (S)-2-methylbutyl, 3-aminopropyl, 2-hexyl, and 4-aminobutyl; **and**

~~(iii) B is optionally selected from the group consisting of~~ cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, oxalan-2-yl, 2-(2R)-bicyclo[2.2.1]-heptyl, oxetan-3-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, 1-pyrrolidinyl and 1-piperidinyl;

A is selected from the group consisting of a bond, CH₂, CH₃CH, CH₂CH₂, and CH₂CH₂CH₂;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxymethyl, amino, aminomethyl, cyano, methyl, trifluoromethyl, fluoro, and chloro;

R² is Z⁰-Q;

Z⁰ is selected from the group consisting of a bond, O, S, NH, and OCH₂;

Q is selected from the group consisting of

3-amidocarbonyl-5-aminophenyl, 3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-benzylamidosulfonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl,

3-amino-5-(N-ethylamidocarbonyl)phenyl,
3-amino-5-(N-isopropylamidocarbonyl)phenyl,
3-amino-5-(N-propylamidocarbonyl)phenyl,
3-amino-5-(N-isobutylamidocarbonyl)phenyl,
3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl,
3-amino-5-(N-cyclobutylamidocarbonyl)phenyl,
3-amino-5-(N-cyclopentylamidocarbonyl)phenyl,
3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 5-amino-2-fluorophenyl,
3-amino-5-hydroxymethylphenyl, 5-amino-3-methoxycarbonylphenyl,
3-amidinophenyl, 3-amino-2-methylphenyl, 5-amino-2-methylthiophenyl,
3-aminophenyl, 3-carboxyphenyl, 3-carboxy-5-aminophenyl,
3-carboxy-5-hydroxyphenyl, 3-carboxymethyl-5-aminophenyl,
3-carboxymethyl-5-hydroxyphenyl, 3-carboxymethylphenyl, 3-chlorophenyl, 2-chlorophenyl, 3-cyanophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl,
2-fluorophenyl, 3-fluorophenyl, 2,5-difluorophenyl, 2-hydroxyphenyl,
3-hydroxyphenyl, 3-methanesulfonylaminophenyl, 2-methoxyphenyl,
3-methoxyphenyl, 3-methoxyaminophenyl, 3-methoxycarbonylphenyl,
2-methylaminophenyl, 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-trifluoromethylphenyl, 2-trifluoromethylphenyl, 5-amino-2-thienyl, 5-amino-3-thienyl,
3-bromo-2-thienyl, 3-pyridyl, 4-pyridyl, 2-thienyl, and 3-thienyl, with the proviso that Q is other than a phenyl or a substituted phenyl when Z⁰ is a bond;

Y^{AT} is Q^b-Q^s;

Q^s is selected from the group consisting of:

[CH(benzoyl)](CH₂)_b, [CH(2-pyridylcarbonyl)](CH₂)_b,
[CH(3-pyridylcarbonyl)](CH₂)_b, [CH(4-pyridylcarbonyl)](CH₂)_b,
[CH(2-thienylcarbonyl)](CH₂)_b, [CH(3-thienylcarbonyl)](CH₂)_b,
[CH(2-thiazolylcarbonyl)](CH₂)_b, [CH(4-thiazolylcarbonyl)](CH₂)_b,
and [CH(5-thiazolylcarbonyl)](CH₂)_b, wherein b is an integer selected from 1 through 3, with the proviso that said benzoyl and said heteroaroyls are optionally substituted with one or more substituents selected from the group consisting of R¹⁶, R¹⁷, R¹⁸, and R¹⁹ with the proviso that R¹⁷ and R¹⁸ are optionally substituted at a carbon selected from other than the meta and para carbons relative to the carbonyl of the benzoyl or the heteroaroyl, and that said benzoyl or said heteroaroyl are

bonded to the carbon directly bonded to amide nitrogen of the 1-(amidocarbonylmethylene) group;

R¹⁶ and R¹⁹ are independently selected from the group consisting of hydrido, amidino, amino, aminomethyl, methoxy, methylamino, hydroxy, hydroxymethyl, fluoro, chloro, and cyano;

R¹⁷ and R¹⁸ are independently selected from the group consisting of hydrido, fluoro, chloro, hydroxy, hydroxymethyl, amino, carboxy, and cyano; Q^b is N(R²⁶)C(NR²⁵)N(R²³)(R²⁴); **and**

R²³, R²⁴, R²⁵, and R²⁶ are independently hydrido or methyl.

36. (currently amended): The compound as recited in **[[Claim]] claim 35** or a pharmaceutically acceptable salt thereof, wherein;

B is selected from the group consisting of:

(i) 3-aminophenyl, 3-amidinophenyl, 4-amidinophenyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-dichlorophenyl, 2-fluorophenyl, 4-methylphenyl, phenyl, 2-imidazolyl, 3-pyridyl, 4-pyridyl, and 3-trifluoromethyl-2-pyridyl;

(ii) ~~B is optionally selected from the group consisting of~~ hydrido, ethyl, 2-propenyl, 2-propynyl, propyl, isopropyl, butyl, 2-butyl, (R)-2-butyl, (S)-2-butyl, tert-butyl, isobutyl, 1-pentyl, 3-pentyl, 2-methylbutyl, 2,2,2-trifluoroethyl, 6-amidocarbonylhexyl, 4-methyl-2-pentyl, 3-hydroxypropyl, 1-methoxy-2-propyl, 2-methoxyethyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 2-dimethylaminopropyl, 2-cyanoethyl, 6-hydroxyhexyl, 2-hydroxyethyl, 2-amidinoethyl, 2-guanidinoethyl, 3-guanidinopropyl, 4-guanidinobutyl, 3-hydroxypropyl, 4-hydroxybutyl, 6-cyanoethyl, 2-dimethylaminoethyl, 3-methylbutyl, 2-methylbutyl, (S)-2-methylbutyl, 3-aminopropyl, 2-hexyl, and 4-aminobutyl; **and**

(iii) ~~B is optionally selected from the group consisting of~~ cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, oxalan-2-yl, 2-(2R)-bicyclo[2.2.1]-heptyl, oxetan-3-yl, azetidin-1-yl, azetidin-2-yl, azetidin-3-yl, and 1-piperidinyl;

A is selected from the group consisting of a bond, CH₂, CH₂CH₂ and CH₂CH₂CH₂;

M is N or R¹-C;

R¹ is selected from the group consisting of hydrido, hydroxy, hydroxymethyl, amino, aminomethyl, cyano, methyl, trifluoromethyl, and fluoro;

R² is Z⁰-Q;


Z⁰ is selected from the group consisting of a bond, O, S, and NH;

Q is selected from the group consisting of

3-amidocarbonyl-5-aminophenyl, 3-amino-5-(N-benzylamidocarbonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(3-fluorobenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-(1-methyl-1-phenylethyl)amidocarbonyl)phenyl, 3-amino-5-(N-benzylamidosulfonyl)phenyl, 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, 3-amino-5-(N-ethylamidocarbonyl)phenyl, 3-amino-5-(N-isopropylamidocarbonyl)phenyl, 3-amino-5-(N-propylamidocarbonyl)phenyl, 3-amino-5-(N-isobutylamidocarbonyl)phenyl, 3-amino-5-(N-(2-butyl)amidocarbonyl)phenyl, 3-amino-5-(N-cyclobutylamidocarbonyl)phenyl, 3-amino-5-(N-cyclopentylamidocarbonyl)phenyl, 3-amino-5-(N-cyclohexylamidocarbonyl)phenyl, 3-aminophenyl, 3-carboxy-5-aminophenyl, 3-chlorophenyl, 3,5-diaminophenyl, 3-dimethylaminophenyl, 3-hydroxyphenyl, 3-methanesulfonylaminophenyl, 3-methylaminophenyl, 2-methylphenyl, 3-methylphenyl, phenyl, 3-trifluoroacetamidophenyl, 3-bromo-2-thienyl, 2-thienyl, and 3-thienyl, with the proviso that Q is other than a phenyl or a substituted phenyl when Z⁰ is a bond; **and**

Y^{AT} is selected from the group consisting of 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, 5-guanidino-1-oxo-1-(4-thiazolyl)-2-pentyl, 5-guanidino-1-oxo-1-(5-thiazolyl)-2-pentyl, 5-guanidino-1-oxo-1-(4-amino-2-thiazolyl)-2-pentyl, and 5-guanidino-1-oxo-1-phenyl-2-pentyl.

37. (currently amended): A compound as recited in [[Claim]] **claim** 33 where said compound is selected from the group of the Formula:



or a pharmaceutically acceptable salt thereof, wherein:

R^2 is 3-aminophenoxy, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3,5-diaminophenoxy, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-carboxy-5-aminophenoxy, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenoxy, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3,5-diaminophenoxy, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-carboxy-5-aminophenoxy, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenoxy, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3,5-diaminophenoxy, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-carboxy-5-aminophenoxy, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenoxy, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH.

R^2 is 3-aminophenylthio, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3,5-diaminophenylthio, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-carboxy-5-aminophenylthio, B is phenyl, A is CH_2CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenylthio, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3,5-diaminophenylthio, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-carboxy-5-aminophenylthio, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenylthio, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3,5-diaminophenylthio, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-carboxy-5-aminophenylthio, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenylthio, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3,5-diamino-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-carboxy-5-amino-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3,5-diamino-2-thienyl, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-carboxy-5-amino-2-thienyl, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3,5-diamino-2-thienyl, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-carboxy-5-amino-2-thienyl, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R² is 3-aminophenoxy, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diaminophenoxy, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-aminophenoxy, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenoxy, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diaminophenoxy, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-aminophenoxy, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenoxy, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diaminophenoxy, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-aminophenoxy, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenoxy, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N.

R² is 3-aminophenylthio, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diaminophenylthio, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-aminophenylthio, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenylthio, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diaminophenylthio, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-aminophenylthio, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenylthio, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diaminophenylthio, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-aminophenylthio, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenylthio, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diamino-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-amino-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is phenyl, A is CH₂CH₂, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diamino-2-thienyl, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-amino-2-thienyl, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is isopropyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3,5-diamino-2-thienyl, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N;

R² is 3-carboxy-5-amino-2-thienyl, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N; or

R² is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is cyclobutyl, A is single bond, Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N.

38. (currently amended): A composition for inhibiting thrombotic conditions in blood comprising a compound of ~~any one of Claims~~ claims 8, 16, 24, 32, ~~[[and]]~~ or 37 and a pharmaceutically acceptable carrier.

39. (currently amended): A composition for inhibiting thrombotic conditions in blood comprising a compound of ~~any one of Claims 1 through 7, Claims 9 through 15, Claims 17 through 23, Claims 25 through 31, and Claims 33 through 36~~ claims 1, 9, 17, 25, or 33 and a pharmaceutically acceptable carrier.

40. (currently amended): A method for inhibiting thrombotic conditions in blood comprising adding to blood a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

41. (currently amended): A method for inhibiting formation of blood platelet aggregates in blood comprising adding to blood a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

42. (currently amended): A method for inhibiting thrombus formation in blood comprising adding to blood a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

43. (currently amended): A method for treating or preventing venous thromboembolism and pulmonary embolism in a mammal comprising administering to the mammal a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

44. (currently amended): A method for treating or preventing deep vein thrombosis in a mammal comprising administering to the mammal a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

45. (currently amended): A method for treating or preventing cardiogenic thromboembolism in a mammal comprising administering to the mammal a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

46. (currently amended): A method for treating or preventing thromboembolic stroke in humans and other mammals comprising administering to the mammal a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

47. (currently amended): A method for treating or preventing thrombosis associated with cancer and cancer chemotherapy in humans and other mammals comprising administering to the mammal a therapeutically effective amount of a composition of **[[Claims]] claim 38**.

48. (currently amended): A method for treating or preventing unstable angina in humans and other mammals comprising administering to the mammal a therapeutically effective amount of a composition of any one of ~~[[Claims]]~~ **claim** 38.

49. (currently amended): A method for inhibiting thrombus formation in blood comprising adding to blood a therapeutically effective amount of a compound of any one of ~~Claims 1 through 37~~ **claims 1, 9, 17, 25, or 33** with a therapeutically effective amount of fibrinogen receptor antagonist.

Claim 50 (canceled).

51. (currently amended): A method of treating or preventing a TF VIIA-mediated disorder in a subject by administering a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof, said compound selected from the group consisting of:

2-[3-[2-[3-aminophenyl]-6-chloro-N-[[4-aminoiminomethylphenyl]methyl]-5-[N,N-dimethylhydrazino]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

2-[3-[2-[3-aminophenyl]-6-chloro-5-[N-ethyl-N-methylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

2-[3-[2-[3-aminophenyl]-6-chloro-5-[N,N-diethylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;

2-[3-[2-[3-aminophenyl]-5-[N-(azetidin-1-yl)amino]-6-chloro-N-[[4-aminoiminomethylphenyl]methyl]-4-oxo-1(4H)-pyrimidinyl]]acetamide;


[2-[4-[3-[3-aminophenyl]-N-[[4-aminoiminomethylphenyl]methyl]-6-[N,N-dimethylhydrazino]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide;

2-[4-[3-[3-aminophenyl]-6-[N-ethyl-N-methylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide;

2-[4-[3-[3-aminophenyl]-6-[N,N-diethylhydrazino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide; **and**

2-[4-[3-[3-aminophenyl]-6-[N-(azetidin-1-yl)amino]-N-[[4-aminoiminomethylphenyl]methyl]-5-oxo-1(5H)-1,2,4-triazinyl]]acetamide.]

52. (currently amended): A method of treating or preventing a TF VIIA-mediated disorder in a subject by administering a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof, said compound of the formula:



wherein:

R² is 3-aminophenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is phenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-dimethylaminophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 2-methylphenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is phenyl, B is 3-aminophenyl, A is C(O)NH, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is phenyl, B is 3-amidinophenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-(N-methylamino)phenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-methylsulfonamidophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is phenyl, B is 4-amidinophenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-methylaminophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is phenyl, B is phenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-methylphenyl, B is 4-phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3-aminophenyl, B is phenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CCl;

R^2 is 3-dimethylaminophenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 2-methylphenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is phenyl, B is 3-aminophenyl, A is $C(O)NH$, Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is phenyl, B is 3-amidinophenyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-(N-methylamino)phenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-methylsulfonamidophenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is phenyl, B is 4-amidinophenyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-methylaminophenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is phenyl, B is phenyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-methylphenyl, B is 4-phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-aminophenyl, B is 3-chlorophenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is 3-aminophenyl, B is phenyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is phenyl, B is 3-chlorophenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is 3-dimethylaminophenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is 2-methylphenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is phenyl, B is 3-aminophenyl, A is $C(O)NH$, Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is phenyl, B is 3-amidinophenyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is 3-(N-methylamino)phenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R^2 is 3-methylsulfonamidophenyl, B is phenyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CF;

R² is phenyl, B is 4-amidinophenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CF;
R² is 3-methylaminophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and
M is CF;

R² is phenyl, B is phenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-methylphenyl, B is 4-phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is
CF;

R² is 3-aminophenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl,
and M is N;

R² is 3-aminophenyl, B is phenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is
N;

R² is 3-dimethylaminophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl,
and M is N;

R² is 2-methylphenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is
N;

R² is phenyl, B is 3-aminophenyl, A is C(O)NH, Y⁰ is 4-amidinobenzyl, and M is
N;

R² is phenyl, B is 3-amidinophenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-(N-methylamino)phenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl,
and M is N;

R² is 3-methylsulfonamidophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-
amidinobenzyl, and M is N;

R² is phenyl, B is 4-amidinophenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-methylaminophenyl, B is phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and
M is N;

R² is phenyl, B is phenyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-methylphenyl, B is 4-phenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and
M is N;

R² is 3-amidocarbonyl-5-aminophenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-
amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 3-chlorophenyl, A is
CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 3-chlorophenyl, A
is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amidocarbonyl-5-aminophenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3,5-diaminophenoxy, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-carboxy-2-thienyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2,2,2-trifluoroethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is (S)-2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 5-amino-2-fluorophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 2-methyl-3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is ethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is ethyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propenyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is isopropyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 2-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is (R)-2-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 2-propynyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 3-pentyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is hydrido, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is ethyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 2-methypropyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 2-propyl, A is CH_3CH , Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is propyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 6-amidocarbonylhexyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is tert-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is tert-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 3-hydroxypropyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 2-methylpropyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 3-methoxy-2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is 3-methoxy-2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-methoxy-2-ethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 5-amidino-2-thienylmethyl, and M is CH;

R² is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is CH;

R² is 3-carboxyphenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2,2,2-trifluoroethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is (S)-2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 5-amino-2-fluorophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 2-methyl-3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is ethyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is ethyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is N;

R² is 3-aminophenyl, B is 2-propenyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is N;

R² is 3-aminophenyl, B is isopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is 2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is (R)-2-butyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is 2-propynyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 3-pentyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is hydrido, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is ethyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-methylpropyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-propyl, A is CH_3CH , Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is propyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 6-amidocarbonylhexyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is tert-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is tert-butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 3-hydroxypropyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-methylpropyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is N;

R^2 is 3-aminophenyl, B is butyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 3-methoxy-2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 3-methoxy-2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-methoxy-2-ethyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-propyl, A is single bond, Y^0 is 5-amidino-2-thienylmethyl, and M is N;

R^2 is 3-aminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidino-3-fluorobenzyl, and M is N;

R^2 is 3-carboxyphenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidino-3-fluorobenzyl, and M is CH;

R^2 is 3-amidocarbonyl-5-aminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3,5-diaminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-amino-5-carboxyphenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-amidocarbonyl-5-aminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3,5-diaminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amino-5-carboxyphenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amino-5-carbomethoxyphenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-amidocarbonyl-5-aminophenyl, B is 2-propyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3,5-diaminophenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3-amino-5-carboxyphenyl, B is 2-propyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R² is 3-aminophenyl, B is cyclopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclopropyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclopentyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 5-amino-2-thienyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclopropyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is 2-(2R)-bicyclo[2.2.1]-heptyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-aminophenyl, B is cyclopentyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is cyclohexyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 2-hydroxyphenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is phenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-thienyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 2,6-dichlorophenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CH;

R^2 is 3-aminophenyl, B is cyclopropyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclopropyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidino-3-fluorobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclopentyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 5-amino-2-thienyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclopropyl, A is CH_2 , Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is 2-(2R)-bicyclo[2.2.1]-heptyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclopentyl, A is single bond, Y^0 is 4-amidino-2-fluorobenzyl, and M is N;

R^2 is 3-aminophenyl, B is cyclohexyl, A is CH_2CH_2 , Y^0 is 4-amidinobenzyl, and M is N;

R² is 2-hydroxyphenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-thienyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 2,6-dichlorophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-aminophenyl, B is cyclopropyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclopropyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidino-3-fluorobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclopentyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 5-amino-2-thienyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclopropyl, A is CH₂, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-aminophenyl, B is 2-(2R)-bicyclo[2.2.1]-heptyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclopentyl, A is single bond, Y⁰ is 4-amidino-2-fluorobenzyl, and M is CF;

R² is 3-aminophenyl, B is cyclohexyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 2-hydroxyphenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-thienyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 2,6-dichlorophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CF;

R² is 3-amidocarbonyl-5-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3,5-diaminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amino-5-carboxyphenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is N;

R² is 3-amidocarbonyl-5-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3,5-diaminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amino-5-carboxyphenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CH;

R² is 3-amidocarbonyl-5-aminophenyl, B is cyclobutyl, A is single bond, Y⁰ is 4-amidinobenzyl, and M is CCl;

R^2 is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl;


R^2 is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl;

R^2 is 3,5-diaminophenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl; or

R^2 is 3-amino-5-carboxyphenyl, B is cyclobutyl, A is single bond, Y^0 is 4-amidinobenzyl, and M is CCl.

53. (currently amended): A method of treating or preventing a TF VIIA-mediated disorder in a subject by administering a therapeutically effective amount of a compound or a pharmaceutically acceptable salt thereof, said compound of the formula:



wherein;


R^2 is 3-aminophenyl, B is phenyl, A is CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CH;

R^2 is 3-aminophenyl, B is phenyl, A is CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CF;

R^2 is 3-aminophenyl, B is phenyl, A is CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is CCl; or


R^2 is 3-aminophenyl, B is phenyl, A is CH_2 , Y^{AT} is 5-guanidino-1-oxo-1-(2-thiazolyl)-2-pentyl, and M is N.

54. (new): A compound of the Formula:



or a pharmaceutically acceptable salt thereof, wherein:

B is the Formula:



R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b ;

A is selected from the group consisting of single covalent bond and $(CH(R^{15}))_{pa}-(W^7)_r$ wherein r is an integer selected from 0 through 1, pa is an integer selected from 0 through 3, and W^7 is $N(R^7)$;

R^7 is selected from the group consisting of hydrido and alkyl;

R^{15} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is N or R^1-C ;

R^1 is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

Q is selected from the group consisting of aryl and heteroaryl wherein (a) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (b) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (c) a ring carbon or nitrogen, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (d) a ring carbon or nitrogen, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², and (e) a ring carbon or nitrogen, if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R¹⁰ and R¹², respectively, is optionally substituted by R¹¹;

R⁹, R¹¹, and R¹³ are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkoxy, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, monoalkyl amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboxy, carboxamido, and cyano;

R¹⁰ and R¹² are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, alkoxy, alkoxyamino, aminoalkyl, hydroxy, amino, alkylamino, alkylsulfonamido, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, hydroxyalkyl, aminoalkyl, halo, haloalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxyamido, and cyano;

Y⁰ is amidinoaralkyl or amidinoheteroaralkyl;

Q^b is selected from the group consisting of hydrido, NR²⁰R²¹, N(R²⁶)C(NR²⁵)N(R²³)(R²⁴), and C(NR²⁵)NR²³R²⁴; and

R²⁰, R²¹, R²³, R²⁴, R²⁵, and R²⁶ are independently selected from the group consisting of hydrido and alkyl.


55. (new): The compound of claim 54 wherein B is phenyl or phenyl substituted by hydrido, halo, amidino, or hydroxy.

56. (new): The compound of claim 55 wherein A is CH₂ or CH₂CH₂ and M is CH or C-halo.

57. (new): The compound of claim 56 wherein B is phenyl, chlorophenyl, or


amidinophenyl.

58. (new): The compound of claim 57 wherein Q is



and R¹⁰ and R¹² are as defined in claim 54.

59. (new): The compound of claim 54 wherein Q is



and R¹⁰ and R¹² are as defined in claim 54.

60. (new): The compound of claim 54 wherein Q is 3-amino-5-carboxy-2-thienyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

61. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

62. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)-2-thienyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-

amidinobenzyl, and M is CH or CCl.

63. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)-2-thienyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

64. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)-2-thienyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

65. (new): The compound of claim 54 wherein Q is 3-amidocarbonyl-5-aminophenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH.


66. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH.

67. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH.

68. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH.

69. (new): The compound of claim 54 wherein Q is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 3-chlorophenyl, A is CH₂CH₂, Y⁰ is 4-amidinobenzyl, and M is CH.

70. (new):
A compound of the
Formula:



or a pharmaceutically acceptable salt thereof, wherein:

B is selected from the group consisting of hydrido, C2-C8 alkyl, C3-C8 alkenyl, C3-C8 alkynyl, and C2-C8 haloalkyl, wherein each member of group B is optionally substituted at any carbon up to and including 6 atoms from the point of attachment of B to A with one or more of the group consisting of R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} ;

R^{32} , R^{33} , R^{34} , R^{35} , and R^{36} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboalkoxy, carboxy, carboxamido, cyano, and Q^b ;

A is selected from the group consisting of single covalent bond and $(CH(R^{15}))_{pa}-(W^7)_r$, wherein rr is an integer selected from 0 through 1, pa is an integer selected from 0 through 3, and W^7 is $N(R^7)$;

R^7 is selected from the group consisting of hydrido and alkyl;

R^{15} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is N or R^1-C

R^1 is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

Q is selected from the group consisting of aryl and heteroaryl wherein (a) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon or nitrogen, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon or nitrogen, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon or nitrogen, if present, in the gamma position relative to the ring carbon at the point of

attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11} ;

R^9 , R^{11} , and R^{13} are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkoxy, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, monoalkyl amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboxy, carboxamido, and cyano;

R^{10} and R^{12} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, alkoxy, alkoxyamino, aminoalkyl, hydroxy, amino, alkylamino, alkylsulfonamido, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, hydroxyalkyl, aminoalkyl, halo, haloalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxyamido, and cyano;

Y^0 is amidinoaralkyl or amidinoheteroaralkyl;

Q^b is selected from the group consisting of hydrido, $NR^{20}R^{21}$, $N(R^{26})C(NR^{25})N(R^{23})(R^{24})$, and $C(NR^{25})NR^{23}R^{24}$; and


R^{20} , R^{21} , R^{23} , R^{24} , R^{25} , and R^{26} are independently selected from the group consisting of hydrido and alkyl.

71. (new): The compound of claim 70 wherein B is hydrido, C2-C8 alkyl, C3-C8 alkenyl, C3-C8 alkynyl, or C2-C8 haloalkyl.

72. (new): The compound of claim 71 wherein A is a bond and M is CH or CCl.


73. (new): The compound of claim 72 wherein B is C2-C8 alkyl.

74. (new): The compound of claim 73 wherein Q is




and R^{10} and R^{12} are as defined in claim 70.

75. (new): The compound of claim 70 wherein Q is




and R^{10} and R^{12} are as defined in claim 70.

76. (new): The compound of claim 70 wherein the compound is



77. (new): The compound of claim 70 wherein the compound is



78. (new): The compound of claim 70 wherein Q is 3-amino-5-carboxy-2-thienyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

79. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

80. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)-2-thienyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

81. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)-2-thienyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

82. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)-2-thienyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

83. (new): The compound of claim 70 wherein Q is 3,5-diamino-2-thienyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

84. (new): The compound of claim 70 wherein Q is 3-amidocarbonyl-5-aminophenyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

85. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.


86. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

87. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

88. (new): The compound of claim 70 wherein Q is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

89. (new): The compound of claim 70 wherein Q is 3,5-diaminophenyl, B is 2-propyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

90. (new): A compound of the Formula:



or a pharmaceutically acceptable salt thereof, wherein:

B is selected from the group consisting of C3-C7 cycloalkyl and C4-heterocyclyl, wherein (a) each ring carbon is optionally substituted with R³³, (b) a ring carbon, other than the ring carbon at the point of attachment, is optionally substituted with oxo provided that no more than one ring carbon is substituted by oxo at the same time, (c) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R⁹, (d) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R¹³, (e) a ring carbon or nitrogen, if present, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R⁹, is optionally substituted by R¹⁰, (f) a ring carbon or nitrogen, if present, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹³, is optionally substituted by R¹², (g) a ring carbon or nitrogen, if present, in a first gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R¹⁰, is optionally substituted by R¹¹, and (h) a ring carbon or nitrogen, if present, in a second gamma position relative to the

carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{12} , is optionally substituted by R^{33} ;

R^9 , R^{11} , and R^{13} are independently selected from the group consisting of hydrido, hydroxy, amino, amidino, guanidino, alkylamino, alkylthio, alkoxy, alkylsulfinyl, alkylsulfonyl, amidosulfonyl, monoalkyl amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboxy, carboxamido, and cyano;

R^{10} and R^{12} are independently selected from the group consisting of hydrido, acetamido, haloacetamido, amidino, guanidino, alkyl, alkoxy, alkoxyamino, aminoalkyl, hydroxy, amino, alkylamino, alkylsulfonamido, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, hydroxyalkyl, aminoalkyl, halo, haloalkyl, carboalkoxy, carboxy, carboxyalkyl, carboxyamido, and cyano;

R^{33} and R^{34} are independently selected from the group consisting of hydrido, amidino, guanidino, alkoxy, hydroxy, amino, alkoxyamino, alkylamino, alkylthio, amidosulfonyl, monoalkyl amidosulfonyl, dialkyl amidosulfonyl, alkyl, halo, haloalkyl, haloalkoxy, hydroxyalkyl, carboalkoxy, carboxy, carboxamido, and cyano;

R^{33} is optionally Q^b ;

A is selected from the group consisting of single covalent bond and $(CH(R^{15}))_{pa}-(W^7)_r$ wherein rr is an integer selected from 0 through 1, pa is an integer selected from 0 through 3, and W^7 is $N(R^7)$;

R^7 is selected from the group consisting of hydrido, hydroxy and alkyl;

R^{15} is selected from the group consisting of hydrido, halo, alkyl, and haloalkyl;

M is R^1-C .

R^1 is selected from the group consisting of hydrido, hydroxy, hydroxyamino, amidino, amino, cyano, hydroxyalkyl, alkoxy, alkyl, alkylamino, aminoalkyl, alkylthio, alkoxyamino, haloalkyl, haloalkoxy, and halo;

Q is selected from the group consisting of aryl and heteroaryl wherein (a) a ring carbon or nitrogen in a first alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^9 , (b) a ring carbon or nitrogen in a second alpha position relative to the ring carbon at the point of attachment is optionally substituted by R^{13} , (c) a ring carbon or nitrogen, in a first beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^9 , is optionally substituted by R^{10} , (d) a ring carbon or nitrogen, in a second beta position relative to the ring carbon at the point of attachment and in an alpha position relative to the ring atom optionally substituted by R^{13} , is optionally substituted by R^{12} , and (e) a ring carbon or nitrogen,

if present, in the gamma position relative to the ring carbon at the point of attachment and in an alpha position relative to each of the ring atoms optionally substituted by R^{10} and R^{12} , respectively, is optionally substituted by R^{11} ;

Y^0 is amidinoaralkyl or amidinoheteroaralkyl;

Q^b is selected from the group consisting of hydrido, $NR^{20}R^{21}$, and $C(NR^{25})NR^{23}R^{24}$; and


R^{20} , R^{21} , R^{23} , R^{24} , and R^{25} are independently selected from the group consisting of hydrido and alkyl.

91. (new): The compound of claim 90 wherein B is C3-C7 cycloalkyl or C4-heterocyclyl.

92. (new): The compound of claim 91 wherein A is a bond and M is CH or C-halo.


93. (new): The compound of claim 92 wherein B is C3-C7 cycloalkyl.

94. (new): The compound of claim 93 wherein Q is



and R^{10} and R^{12} are as defined in claim 90.

95. (new): The compound of claim 90 wherein R^2 is



and R¹⁰ and R¹² are as defined in claim 90.

96. (new): The compound of claim 90 wherein Q is 3-amino-5-carboxy-2-thienyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

97. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-benzylamidocarbonyl)-2-thienyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

98. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)-2-thienyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

99. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)-2-thienyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

100. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)-2-thienyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

101. (new): The compound of claim 90 wherein Q is 3,5-diamino-2-thienyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

102. (new): The compound of claim 90 wherein Q is 3-amidocarbonyl-5-aminophenyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

103. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-benzylamidocarbonyl)phenyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

104. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

105. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-(2-chlorobenzyl)amidosulfonyl)phenyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

106. (new): The compound of claim 90 wherein Q is 3-amino-5-(N-(2-trifluoromethylbenzyl)amidocarbonyl)phenyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.

107. (new): The compound of claim 90 wherein Q is 3,5-diaminophenyl, B is cyclobutyl, A is a single bond, Y⁰ is 4-amidinobenzyl, and M is CH or CCl.